

Datasheet: MCA643R

BATCH NUMBER 172783

Description:	MOUSE ANTI RAT CD44
Specificity:	CD44
Other names:	H-CAM, PGP-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-50
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Rat T cell blasts.
External Database Links	<p>UniProt: P26051 Related reagents</p> <p>Entrez Gene: 25406 Cd44 Related reagents</p>
RRID	AB_566756
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS0/1 myeloma cell line.
Specificity	<p>Mouse anti Rat CD44 antibody, clone OX-50 recognizes the rat CD44 cell surface antigen, also known as Extracellular matrix receptor III, P90 lymphocyte homing/adhesion receptor, HUTCH-I, Hermes antigen, Hyaluronate receptor, Phagocytic glycoprotein 1, PGP-1 or Phagocytic glycoprotein I.</p> <p>CD44 is a 482 amino acid ~85 kDa single pass type I transmembrane glycoprotein, expressed by T cells, B cells, macrophages and thymocytes, with expression being increased following activation.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Stevenson, K.S. <i>et al.</i> (2009) Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations. Stem Cells Dev. 18:1389-98. 2. Jiang, T.S. <i>et al.</i> (2010) Reconstruction of the corneal epithelium with induced marrow mesenchymal stem cells in rats. Mol Vis. 16: 1304-16. 3. Kanellis, J. <i>et al.</i> (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. Nephrol Dial Transplant. 25: 2898-908. 4. Li, S. <i>et al.</i> (2010) Upregulation of CXCR4 favoring neural-like cells migration via AKT activation. Neurosci Res. 67: 293-9. 5. Stephens, L.A. <i>et al.</i> (2004) Phenotypic characterization of regulatory CD4+CD25+ T cells in rats. Int Immunol. 16: 365-75. 6. Rice, C.M. <i>et al.</i> (2010) Multipotent adult progenitor cell isolation and proliferation in cytokine and serum-free medium conditioned by rat B104 cells. Br J Haematol. 148: 441-4. 7. Carulli, D. <i>et al</i> (2006) Composition of perineuronal nets in the adult rat cerebellum and

- the cellular origin of their components. [J Comp Neurol. 494: 559-77.](#)
8. Galtrey, C.M. *et al.* (2008) Distribution and synthesis of extracellular matrix proteoglycans, hyaluronan, link proteins and tenascin-R in the rat spinal cord. [Eur J Neurosci. 27: 1373-90.](#)
 9. Hofmann, N. *et al.* (2002) Increased expression of ICAM-1, VCAM-1, MCP-1, and MIP-1 alpha by spinal perivascular macrophages during experimental allergic encephalomyelitis in rats. [BMC Immunol. 3: 11.](#)
 10. Walther, M. *et al.* (2001) Exogenous antigen containing perivascular phagocytes induce a non-encephalitogenic extravasation of primed lymphocytes. [J Neuroimmunol. 117: 30-42.](#)
 11. Suzuki, A. *et al.* (2006) Localization of CD44 and hyaluronan in the synovial membrane of the rat temporomandibular joint. [Anat Rec A Discov Mol Cell Evol Biol. 288: 646-52.](#)
 12. Goransson, V. *et al.* (2004) Renal hyaluronan accumulation and hyaluronan synthase expression after ischaemia-reperfusion injury in the rat. [Nephrol Dial Transplant. 19: 823-30.](#)
 13. Campbell, N.G. *et al.* (2016) Cell Size Critically Determines Initial Retention of Bone Marrow Mononuclear Cells in the Heart after Intracoronary Injection: Evidence from a Rat Model. [PLoS One. 11 \(7\): e0158232.](#)
 14. Bejar, M.T. *et al.* (2016) Inhibition of Notch rescues the angiogenic potential impaired by cardiovascular risk factors in epicardial adipose stem cells. [FASEB J. 30 \(8\): 2849-59.](#)
 15. Rochefort, G.Y. *et al.* (2006) Multipotential mesenchymal stem cells are mobilized into peripheral blood by hypoxia. [Stem Cells. 24 \(10\): 2202-8.](#)
 16. Redondo, J. *et al.* (2015) Reductions in kinesin expression are associated with nitric oxide-induced axonal damage. [J Neurosci Res. 93 \(6\): 882-92.](#)
 17. Huang, X. *et al.* (2019) MRI Tracking of SPIO- and *Fth1*-Labeled Bone Marrow Mesenchymal Stromal Cell Transplantation for Treatment of Stroke. [Contrast Media Mol Imaging. 2019: 5184105.](#)
 18. Aminzadeh, A. *et al.* (2020) Investigating The Alterations of Oxidative Stress Status, Antioxidant Defense Mechanisms, MAP Kinase and Mitochondrial Apoptotic Pathway in Adipose-Derived Mesenchymal Stem Cells from STZ Diabetic Rats. [Cell J. 22 \(Suppl 1\): 38-48.](#)
 19. Paiva, R.G. *et al.* (2020) Stem cells in end-to-side neurorrhaphy. Experimental study in rats [Acta Cirúrgica Brasileira. 35 \(12\) \[Epub ahead of print\].](#)
 20. Hou, B. *et al.* (2018) Comparison of the Effects of BMSC-derived Schwann Cells and Autologous Schwann Cells on Remyelination Using a Rat Sciatic Nerve Defect Model. [Int J Biol Sci. 14 \(13\): 1910-22.](#)
 21. Porwal, K. *et al.* (2019) Increased bone marrow-specific adipogenesis by clofazimine causes impaired fracture healing, osteopenia and osteonecrosis without extra-skeletal effects in rats. [Toxicol Sci. kfz172.](#)
 22. Chang, H.H. *et al.* (2019) Intrarenal Transplantation of Hypoxic Preconditioned Mesenchymal Stem Cells Improves Glomerulonephritis through Anti-Oxidation, Anti-ER Stress, Anti-Inflammation, Anti-Apoptosis, and Anti-Autophagy. [Antioxidants \(Basel\). 9 \(1\): 2.](#)
 23. Wu, J. *et al.* (2022) Reinforcing the function of bone graft via the Ca-P ceramics dynamic behavior-enhanced osteogenic microenvironment for optimal bone regeneration and reconstruction [Applied Materials Today. 27: 101465.](#)

24. Cheng, Y.H. *et al.* (2022) Intrarenal Arterial Transplantation of Dexmedetomidine Preconditioning Adipose Stem-Cell-Derived Microvesicles Confers Further Therapeutic Potential to Attenuate Renal Ischemia/Reperfusion Injury through miR-122-5p/Erythropoietin/Apoptosis Axis. [Antioxidants \(Basel\). 11\(9\): 1702.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA643R>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M381452:210512'

Printed on 29 Jan 2026