

Datasheet: MCA647P

BATCH NUMBER 151286

Description:	MOUSE ANTI HUMAN IgG (Fc) CH2 DOMAIN:HRP
Specificity:	IgG (Fc) (CH2 DOMAIN)
Format:	HRP
Product Type:	Monoclonal Antibody
Clone:	MK 1 A6
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen (1)	■			1/100 - 1/200
Immunohistology - Paraffin		■		
ELISA	■			1/1000 - 1/10000
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	Reacts with: Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid

Preparation	Purified IgG prepared by affinity chromatography on Prosep A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.01% Thiomersal
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human IgG Polyclonal.
External Database Links	<p>UniProt:</p> <p>P01857 Related reagents</p> <p>P01859 Related reagents</p> <p>P01861 Related reagents</p> <p>P01834 Related reagents</p> <p>P01860 Related reagents</p> <p>P0CG04 Related reagents</p> <p>Entrez Gene:</p> <p>3500 IGHG1 Related reagents</p> <p>3501 IGHG2 Related reagents</p> <p>3502 IGHG3 Related reagents</p> <p>3503 IGHG4 Related reagents</p> <p>3514 IGKC Related reagents</p> <p>28815 IGLV2-14 Related reagents</p>
RRID	AB_321911
Fusion Partners	Spleen cells from BALB/c mouse were fused with cells from the mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Human IgG (Fc) CH2 domain, clone MK 1 A6 recognizes human IgG Fc (all subclasses).</p> <p>CH2 and hinge regions have an important role in effector functions of IgG. The epitope detected by clone MK 1 A6 lies within the CH2 domain as determined by haemagglutination and western blotting using IgG heavy chain and myelomas with defined domain deletions.</p>
References	<p>1. Lund, J. <i>et al.</i> (1996) Multiple interactions of IgG with its core oligosaccharide can modulate recognition by complement and human Fc gamma receptor I and influence the synthesis of its oligosaccharide chains. J Immunol. 157 (11): 4963-9.</p> <p>2. Wozniak-Knopp, G. <i>et al.</i> (2010) Introducing antigen-binding sites in structural loops of immunoglobulin constant domains: Fc fragments with engineered HER2/neu-binding sites and antibody properties. Protein Eng Des Sel. 23: 289-97.</p>

3. Raghuraman, S. *et al.* (2012) Spontaneous clearance of chronic hepatitis C virus infection is associated with appearance of neutralizing antibodies and reversal of T-cell exhaustion. [J Infect Dis. 205: 763-71.](#)
4. Hasenhindl, C. *et al.* (2013) Stability assessment on a library scale: a rapid method for the evaluation of the commutability and insertion of residues in C-terminal loops of the CH3 domains of IgG1-Fc. [Protein Eng Des Sel. 26 \(10\): 675-82.](#)
5. Rasti, N. *et al.* (2006) Nonimmune immunoglobulin binding and multiple adhesion characterize Plasmodium falciparum-infected erythrocytes of placental origin. [Proc Natl Acad Sci U S A. 103: 13795-800.](#)
6. Traxlmayr, M.W. *et al.* (2014) Construction of pH-sensitive Her2-binding IgG1-Fc by directed evolution. [Biotechnol J. 9: 1013-22.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10094 available at: <https://www.bio-rad-antibodies.com/SDS/MCA647P>
10094

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[AbGUARD® HRP STABILIZER PLUS \(BUF052A\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052B\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052C\)](#)

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M368759:200529'

Printed on 19 Jan 2024