

# Datasheet: MCA647P

**BATCH NUMBER 1705**

<b>Description:</b>	MOUSE ANTI HUMAN IgG (Fc) CH2 DOMAIN:HRP
<b>Specificity:</b>	IgG (Fc) (CH2 DOMAIN)
<b>Format:</b>	HRP
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MK 1 A6
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen (1)	■			1/100 - 1/200
Immunohistology - Paraffin		■		
ELISA	■			1/1000 - 1/10000
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Rhesus Monkey</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid

<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Prosep A
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.01% Thiomersal
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human IgG Polyclonal.
<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">P01857</a>      <a href="#">Related reagents</a></p> <p><a href="#">P01859</a>      <a href="#">Related reagents</a></p> <p><a href="#">P01861</a>      <a href="#">Related reagents</a></p> <p><a href="#">P01834</a>      <a href="#">Related reagents</a></p> <p><a href="#">P01860</a>      <a href="#">Related reagents</a></p> <p><a href="#">P0CG04</a>      <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">3500</a>    IGHG1      <a href="#">Related reagents</a></p> <p><a href="#">3501</a>    IGHG2      <a href="#">Related reagents</a></p> <p><a href="#">3502</a>    IGHG3      <a href="#">Related reagents</a></p> <p><a href="#">3503</a>    IGHG4      <a href="#">Related reagents</a></p> <p><a href="#">3514</a>    IGKC      <a href="#">Related reagents</a></p> <p><a href="#">28815</a>    IGLV2-14   <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_321911
<b>Fusion Partners</b>	Spleen cells from BALB/c mouse were fused with cells from the mouse NS1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human IgG (Fc) CH2 domain, clone MK 1 A6</b> recognizes human IgG Fc (all subclasses).</p> <p>CH2 and hinge regions have an important role in effector functions of IgG. The epitope detected by clone MK 1 A6 lies within the CH2 domain as determined by haemagglutination and western blotting using IgG heavy chain and myelomas with defined domain deletions.</p>
<b>References</b>	<p>1. Lund, J. <i>et al.</i> (1996) Multiple interactions of IgG with its core oligosaccharide can modulate recognition by complement and human Fc gamma receptor I and influence the synthesis of its oligosaccharide chains. <a href="#">J Immunol. 157 (11): 4963-9.</a></p> <p>2. Wozniak-Knopp, G. <i>et al.</i> (2010) Introducing antigen-binding sites in structural loops of immunoglobulin constant domains: Fc fragments with engineered HER2/neu-binding sites and antibody properties. <a href="#">Protein Eng Des Sel. 23: 289-97.</a></p>

3. Raghuraman, S. *et al.* (2012) Spontaneous clearance of chronic hepatitis C virus infection is associated with appearance of neutralizing antibodies and reversal of T-cell exhaustion. [J Infect Dis. 205: 763-71.](#)
4. Hasenhindl, C. *et al.* (2013) Stability assessment on a library scale: a rapid method for the evaluation of the commutability and insertion of residues in C-terminal loops of the CH3 domains of IgG1-Fc. [Protein Eng Des Sel. 26 \(10\): 675-82.](#)
5. Rasti, N. *et al.* (2006) Nonimmune immunoglobulin binding and multiple adhesion characterize Plasmodium falciparum-infected erythrocytes of placental origin. [Proc Natl Acad Sci U S A. 103: 13795-800.](#)
6. Traxlmayr, M.W. *et al.* (2014) Construction of pH-sensitive Her2-binding IgG1-Fc by directed evolution. [Biotechnol J. 9: 1013-22.](#)

#### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

#### Guarantee

12 months from date of despatch

#### Health And Safety Information

Material Safety Datasheet documentation #10094 available at: <https://www.bio-rad-antibodies.com/SDS/MCA647P>  
10094

#### Regulatory

For research purposes only

## Related Products

### Recommended Useful Reagents

[AbGUARD® HRP STABILIZER PLUS \(BUF052A\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052B\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052C\)](#)

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M368759:200529'

Printed on 19 Jan 2024