

Datasheet: MCA649G

BATCH NUMBER 160517

Description:	RAT ANTI DINITROPHENOL
Specificity:	DINITROPHENOL
Other names:	DNP
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LO-DNP-30
Isotype:	IgE
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			2.5 ug/ml (as detection antibody)
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Chemical
Product Form	Purified IgE - liquid
Preparation	Purified IgE prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide

Approx. Protein Concentrations	IgE concentration 1.0 mg/ml
Immunogen	Nippostrongylus brasiliensis-DNP.
RRID	AB_567293
Specificity	Rat anti Dinitrophenol antibody, clone LO-DNP-30 recognizes the dinitrophenyl (DNP) hapten.
References	<ol style="list-style-type: none"> 1. Pivniouk, V.I. <i>et al.</i> (1999) SLP-76 deficiency impairs signaling via the high-affinity IgE receptor in mast cells. J Clin Invest. 103 (12): 1737-43. 2. Gilchrist, M. <i>et al.</i> (2010) A key role for ATF3 in regulating mast cell survival and mediator release. Blood. 115: 4734-41. 3. Minas, I.S. <i>et al.</i> (2012) Physiological and proteomic approaches to address the active role of ozone in kiwifruit post-harvest ripening. J Exp Bot. 63 (7): 2449-64. 4. Okragly, A.J. <i>et al.</i> (2018) Human mast cells release the migraine-inducing factor pituitary adenylate cyclase-activating polypeptide (PACAP). Cephalalgia. 38 (9): 1564-74. 5. Debrand, E. <i>et al.</i> (2019) An intergenic non-coding RNA promoter required for histone modifications in the human β-globin chromatin domain. PLoS One. 14 (8): e0217532.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA649G 10040
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382466:210513'

Printed on 18 Jan 2024