

Datasheet: MCA649G

Description:	RAT ANTI DINITROPHENOL
Specificity:	DINITROPHENOL
Other names:	DNP
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LO-DNP-30
Isotype:	IgE
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			2.5 ug/ml (as detection antibody)
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Chemical
Product Form	Purified IgE - liquid
Preparation	Purified IgE prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide
Approx. Protein	IgE concentration 1.0 mg/ml

Concentrations

Immunogen Nippostrongylus brasiliensis-DNP.

RRID AB_567293

Specificity **Rat anti Dinitrophenol antibody, clone LO-DNP-30** recognizes the dinitrophenyl (DNP) hapten.

References

1. Pivniouk, V.I. *et al.* (1999) SLP-76 deficiency impairs signaling via the high-affinity IgE receptor in mast cells. [J Clin Invest. 103 \(12\): 1737-43.](#)
2. Gilchrist, M. *et al.* (2010) A key role for ATF3 in regulating mast cell survival and mediator release. [Blood. 115: 4734-41.](#)
3. Minas, I.S. *et al.* (2012) Physiological and proteomic approaches to address the active role of ozone in kiwifruit post-harvest ripening. [J Exp Bot. 63 \(7\): 2449-64.](#)
4. Okragly, A.J. *et al.* (2018) Human mast cells release the migraine-inducing factor pituitary adenylate cyclase-activating polypeptide (PACAP). [Cephalalgia. 38 \(9\): 1564-74.](#)
5. Debrand, E. *et al.* (2019) An intergenic non-coding RNA promoter required for histone modifications in the human β -globin chromatin domain. [PLoS One. 14 \(8\): e0217532.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA649G>
10040

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382466:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)