

Datasheet: MCA691AMO

BATCH NUMBER 148938

Description:	MOUSE IgG2b NEGATIVE CONTROL: Amethyst Orange
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	Amethyst Orange
Product Type:	Negative/Isotype Control
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

Target Species Negative Control

Product Form Purified IgG conjugated to Amethyst Orange - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Amethyst Orange	405	540

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide (NaN₃)
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Specificity **Mouse IgG2b Negative Control** is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative

control in this species.

Test results show that MCA691C is also suitable for use as a negative control with bovine, ovine, porcine, canine and guinea-pig tissues.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.
* This antibody should be used at the same concentration as the test antibody.

References

1. Grant, A.J. *et al.* (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. [Am J Pathol. 160 \(4\): 1445-55.](#)
2. Zheng, X. *et al.* (2002) Interleukin-3, but not granulocyte-macrophage colony-stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. [Immunology. 107 \(3\): 306-15.](#)
3. Dalli, J. *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
4. Kapetanovic, R. *et al.* (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
5. Shoham, T. *et al.* (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. [J Histochem Cytochem. 49 \(10\): 1245-52.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA691AMO>
10041

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M360090:191028'

Printed on 09 Mar 2024
