

Datasheet: MCA691APC

Description:	MOUSE IgG2b NEGATIVE CONTROL:APC
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	APC
Product Type:	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	100 TESTS

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further							
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.							
			No N	Not Determined	Suggested Dilution			
	Flow Cytometry	-			*			
	Where this antibody h	as not been test	ed for use	in a particular teo	chnique this does not			
	necessarily exclude its use in such procedures. Suggested working dilutions are given as							
	a guide only. *This antibody should be used at the same concentration as the test antibody.							
Target Species	Negative Control							
Product Form	Purified IgG conjugated to Allophycocyanin (APC) - lyophilized							
Reconstitution	Reconstitute with 1.0 ml distilled water Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.							
Max Ex/Em	Fluorophore	Excitation Max	(nm) Em	ission Max (nm)				
	APC	650		661				
Preparation	Purified IgG prepared by ion exchange chromatography from tissue culture supernatant							
Buffer Solution	Phosphate buffered sa	aline						
Preservative	0.09% Sodium Azide							
Stabilisers	1% Bovine Serum Albumin							
	5% Sucrose							

RRID	AB_322311
Specificity	Mouse IgG2b Negative Control is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.
	Mouse IgG2b Negative Control is also suitable for use as a negative control with bovine, ovine, porcine, canine and guinea-pig tissues.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or cells or 100ul whole blood
References	 Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. <u>J Histochem Cytochem. 49 (10): 1245-52.</u> Zheng, X. <i>et al.</i> (2002) Interleukin-3, but not granulocyte-macrophage colony- stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. <u>Immunology. 107 (3): 306-15.</u> Grant, A.J. <i>et al.</i> (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. <u>Am J Pathol. 160 (4): 1445-55.</u> Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. <u>Blood. 112 (6): 2512-9.</u> Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. <u>J Immunol. 188: 3382-94.</u> Lawrence, J. <i>et al.</i> (2022) Effects of freezing storage on the stability of maternal cellular and humoral immune components in porcine colostrum. <u>Vet Immunol Immunopathol. 254: 110520.</u> Forner, R. <i>et al.</i> (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. <u>PLoS One. 16 (5): e0249366.</u>
Storage	This product is shipped at ambient temperature. Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	6 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA691APC 20487
Regulatory	For research purposes only

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M441223:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint