

Datasheet: MCA691B BATCH NUMBER 150918

Description:	MOUSE IgG2b NEGATIVE CONTROL:Biotin
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	Biotin
Product Type:	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

antibody recognises a rat cell surface marker, and therefore cannot be used as a negative

Specificity	Mouse IgG2b Negative Control is negative on all human cell	ls and cell lines tested. This
RRID	AB_322529	
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml	
Preservative Stabilisers	0.09% Sodium Azide1% Bovine Serum Albumin	
Buffer Solution	Phosphate buffered saline	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	from tissue culture
Product Form	Purified IgG conjugated to Biotin - liquid	
Target Species	Negative Control	

control in this species. Test results show that MCA691C is also suitable for use as a negative control with bovine, ovine, porcine, canine and guinea-pig tissues. Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. *It is suggested that the user adjusts the concentration of this reagent to be the same as the test reagent. References 1. Grant, A.J. et al. (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. Am J Pathol. 160 (4): 1445-55. 2. Zheng, X. et al. (2002) Interleukin-3, but not granulocyte-macrophage colonystimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. Immunology. 107 (3): 306-15. 3. Dalli, J. et al. (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. Blood. 112 (6): 2512-9. 4. Kapetanovic, R. et al. (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. <u>J Immunol. 188: 3382-94.</u> 5. Shoham, T. et al. (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. J Histochem Cytochem. 49 (10): 1245-52. **Storage** Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch

	12 months from date of deepater
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA691B 10041
Regulatory	For research purposes only

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368770:200529'

Printed on 09 Mar 2024

Email: antibody_sales_us@bio-rad.com