

Datasheet: MCA691F BATCH NUMBER 157690

Description:	MOUSE IgG2b NEGATIVE CONTROL:FITC
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	FITC
Product Type:	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

Target Species	Negative Control					
Product Form	Purified IgG conjugate	1 (FITC) - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (ni	m)		
	FITC	490	525			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum					
Approx. Protein Concentrations	IgG concentration 0.1	mg/ml				
RRID	AB_322276					

Specificity

Mouse IgG2b Negative Control is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Test results show that MCA691C is also suitable for use as a negative control with bovine. ovine, porcine, canine and guinea-pig tissues.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

* It is suggested that the user adjusts the concentration of this reagent to be the same as the test reagents.

References

- 1. Grant, A.J. et al. (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. Am J Pathol. 160 (4): 1445-55.
- 2. Zheng, X. et al. (2002) Interleukin-3, but not granulocyte-macrophage colonystimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. Immunology. 107 (3): 306-15.
- 3. Dalli, J. et al. (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. Blood. 112 (6): 2512-9.
- 4. Kapetanovic, R. et al. (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94.
- 5. Shoham, T. et al. (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. J Histochem Cytochem. 49 (10): 1245-52.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:

https://www.bio-rad-antibodies.com/SDS/MCA691F

10041

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368772:200529'

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint