

Datasheet: MCA691PB

| | |
|----------------------|--|
| Description: | MOUSE IgG2b NEGATIVE CONTROL:Pacific Blue® |
| Specificity: | MOUSE IgG2b NEGATIVE CONTROL |
| Format: | Pacific Blue® |
| Product Type: | Negative/Isotype Control |
| Isotype: | IgG2b |
| Quantity: | 100 TESTS/1ml |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ■ | | | * |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

Target Species Negative Control

Product Form Purified IgG conjugated to Pacific Blue® - liquid

| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
|-----------|---------------|---------------------|-------------------|
| | Pacific Blue® | 410 | 455 |

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide
1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.05 mg/ml

Specificity **Mouse IgG2b Negative Control** is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Test results show that MCA691C is also suitable for use as a negative control with bovine, ovine, porcine, canine and guinea-pig tissues.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.
* This antibody should be used at the same concentration as the test antibody.

References

1. Grant, A.J. *et al.* (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. [Am J Pathol. 160 \(4\): 1445-55.](#)
2. Zheng, X. *et al.* (2002) Interleukin-3, but not granulocyte-macrophage colony-stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. [Immunology. 107 \(3\): 306-15.](#)
3. Dalli, J. *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
4. Kapetanovic, R. *et al.* (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
5. Shoham, T. *et al.* (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. [J Histochem Cytochem. 49 \(10\): 1245-52.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

Acknowledgements

The Pacific Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays, and are covered by pending and issued patents.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M335705:181204'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)