

## Datasheet: MCA691PE BATCH NUMBER 162471

Description:	MOUSE IgG2b NEGATIVE CONTROL:RPE
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	RPE
Product Type:	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>				
	rad-antibodies.com/protoc				
	Flow Cytometry	Yes No	Not Determined	Suggested Dilution	
		- ot boon tostad f	or uso in a particular too	brique this does not	
	Where this antibody has not been tested for use in a particular technique this does no				
	necessarily exclude its use in such procedures. Suggested working dilutions are give a guide only.				
	a guide offiy.				
Target Species	Negative Control				
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized				
Reconstitution	Reconstitute with 1 ml distilled water				
Max Ex/Em	FluorophoreExRPE 488nm laser	citation Max (nm) 496	Emission Max (nm) 578		
Preparation	Purified IgG prepared by a supernatant	affinity chromato	graphy on Protein A fron	n tissue culture	
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin				
	5% Sucrose				
RRID	AB_322277				

Specificity	<b>Mouse IgG2b Negative Control</b> is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.			
	Test results show that MCA691C is also suitable for use as a negative control with bovine ovine, porcine, canine and guinea-pig tissues.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.			
References	<ol> <li>Grant, A.J. <i>et al.</i> (2002) Hepatic expression of secondary lymphoid chemokine (CCL21 promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. <u>Am J Pathol. 160 (4): 1445-55.</u></li> <li>Zheng, X. <i>et al.</i> (2002) Interleukin-3, but not granulocyte-macrophage colony-stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. <u>Immunology. 107 (3): 306-15.</u></li> </ol>			
	<ol> <li>Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. <u>Blood. 112 (6): 2512-9.</u></li> <li>Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. <u>J Immunol. 188: 3382-94.</u></li> <li>Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. <u>J Histochem Cytochem. 49 (10): 1245-52.</u></li> </ol>			
Storage	Prior to reconstitution store at +4 <sup>o</sup> C. Following reconstitution store at +4 <sup>o</sup> C. This product should be stored undiluted.			
	DO NOT FREEZE. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.			
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA691PE 20487			
Regulatory	For research purposes only			
orth & South Tel: +1 800 26 merica Fax: +1 919 87 Email: antibod				
To find a batch/lot spec	ific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheet 'M411220:221102'			

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint