

# Datasheet: MCA691PE

### **BATCH NUMBER 1808**

Description:	MOUSE IgG2b NEGATIVE CONTROL:RPE
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	RPE
<b>Product Type:</b>	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	100 TESTS

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	<b>Suggested Dilution</b>
Flow Cytometry				*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

Target Species	Negative Control		
Product Form	Purified IgG conjugate	ed to R. Phycoerythrin	(RPE) - lyophilized
Reconstitution	Reconstitute with 1 m	l distilled water	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	supernatant	by affinity chromatog	raphy on Protein A f
Buffer Solution	Phosphate buffered s	aline	
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum	Albumin	

Specificity	Mouse IgG2b Negative Control is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.  Test results show that MCA691C is also suitable for use as a negative control with bovine,
	ovine, porcine, canine and guinea-pig tissues.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.
References	1. Grant, A.J. <i>et al.</i> (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. <u>Am J Pathol. 160 (4): 1445-55.</u>
	2. Zheng, X. <i>et al.</i> (2002) Interleukin-3, but not granulocyte-macrophage colony-stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. <a href="mailto:lmmunology.107">lmmunology. 107</a> (3): 306-15.
	3. Dalli, J. et al. (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. <u>Blood. 112 (6): 2512-9.</u>
	4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble human
	macrophages in their response to bacterial lipopolysaccharide. <u>J Immunol. 188: 3382-94.</u>
	5. Shoham, T. et al. (2001) Reduced expression of activin A in focal lymphoid
	agglomerates within nasal polyps. <u>J Histochem Cytochem. 49 (10): 1245-52.</u>
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.
	This product should be stored undiluted.
	DO NOT FREEZE. This product is photosensitive and should be protected from light.
	Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA691PE">https://www.bio-rad-antibodies.com/SDS/MCA691PE</a> 20487

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

For research purposes only

Regulatory

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375645:210104'

### Printed on 09 Mar 2024