

Datasheet: MCA691XZ BATCH NUMBER 160106

Description:	MOUSE IgG2b NEGATIVE CONTROL:Preservative Free
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	Preservative Free
Product Type:	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	1 mg

Product Details

Applications	This product has been re	ported to	work in th	e following applicatio	ns. This information is
	derived from testing withi	n our lab	oratories,	peer-reviewed publica	ations or personal
	communications from the	originato	ors. Please	e refer to references in	ndicated for further
	information. For general p	protocol r	ecommen	dations, please visit <u>v</u>	<u>vww.bio-</u>
	rad-antibodies.com/proto	<u>cols</u> .			
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				*
	Immunohistology - Frozen				*
	Immunohistology - Paraffin	-			*
	Immunohistology - Resin	•			*
	Radioimmunoassays				*
	Where this antibody has	not been	tested for	use in a particular teo	chnique this does not
	necessarily exclude its us	se in sucl	h procedur	es. Suggested workir	ng dilutions are given as
	a guide only.				
	*It is recommended that t	the user a	adjusts the	concentration of this	reagent to be the same
	as the test reagents.		,		0
Target Species	Negative Control				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by	affinity cl	hromatogra	aphy on Protein A	
Buffer Solution	Phosphate buffered saline	e			
Preservative Stabilisers	None present				
Approx. Protein Concentrations	lgG concentration 1.0 mg	./ml			

	Test results show that MCA691C is also suitable for use as a negative control with bovir
	ovine, porcine, canine and guinea-pig tissues.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	 Grant, A.J. <i>et al.</i> (2002) Hepatic expression of secondary lymphoid chemokine (CCL2 promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. <u>Am J Pathol. 160 (4): 1445-55.</u> Zheng, X. <i>et al.</i> (2002) Interleukin-3, but not granulocyte-macrophage colony-stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. <u>Immunology. 107 (3): 306-15.</u> Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of
Storage	 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. <u>J Immunol. 188: 3382-94</u> 5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. <u>J Histochem Cytochem. 49 (10): 1245-52</u>. Store at -20°C only.
Storage	 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. <u>J Immunol. 188: 3382-94</u> 5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. <u>J Histochem Cytochem. 49 (10): 1245-52.</u>
Storage	 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94 5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. J Histochem Cytochem. 49 (10): 1245-52. Store at -20°C only. This product should be stored undiluted.
	 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94 5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. J Histochem Cytochem. 49 (10): 1245-52. Store at -20°C only. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we
Guarantee Health And Safety	 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94 5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. J Histochem Cytochem. 49 (10): 1245-52. Store at -20°C only. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94 5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. J Histochem Cytochem. 49 (10): 1245-52. Store at -20°C only. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368778:200529'

Printed on 09 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint