

## Datasheet: MCA691XZ BATCH NUMBER 162013

Description:	MOUSE IgG2b NEGATIVE CONTROL:Preservative Free
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	Preservative Free
Product Type:	Negative/Isotype Control
Isotype:	lgG2b
Quantity:	1 mg

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			*
Immunohistology - Frozen	•			*
Immunohistology - Paraffin	•			*
Immunohistology - Resin	•			*
Radioimmunoassays	•			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

\*It is recommended that the user adjusts the concentration of this reagent to be the same as the test reagents.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Regulatory	For research purposes only
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA691XZ">https://www.bio-rad-antibodies.com/SDS/MCA691XZ</a> 10162
Guarantee	12 months from date of despatch
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
	This product should be stored undiluted.
Storage	Store at -20°C only.
	stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-kappaB-dependent and -independent pathways. <a href="Immunology.107">Immunology. 107 (3): 306-15</a> .  3. Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. <a href="Blood.112">Blood. 112 (6): 2512-9</a> .  4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble huma macrophages in their response to bacterial lipopolysaccharide. <a href="Jmmunol.188:3382-9">Jmmunol. 188: 3382-9</a> .  5. Shoham, T. <i>et al.</i> (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. <a href="JHistochem Cytochem.49">JHistochem Cytochem.49 (10): 1245-52</a> .
References	1. Grant, A.J. <i>et al.</i> (2002) Hepatic expression of secondary lymphoid chemokine (CCL2 promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. Am J Pathol. 160 (4): 1445-55.  2. Zheng, X. <i>et al.</i> (2002) Interleukin-3, but not granulocyte-macrophage colony-
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
	Test results show that MCA691C is also suitable for use as a negative control with bovi ovine, porcine, canine and guinea-pig tissues.
Specificity	<b>Mouse IgG2b Negative Control</b> is negative on all human cells and cell lines tested. The antibody recognises a rat cell surface marker, and therefore cannot be used as a negat control in this species.

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368778:200529'

Printed on 09 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint