

Datasheet: MCA699

BATCH NUMBER 148954

Description:	RAT ANTI HUMAN CD49f
Specificity:	CD49f
Other names:	INTEGRIN ALPHA 6 CHAIN, VLA-6
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	NKI-GoH3
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Mouse, Dog, Pig, Cynomolgus monkey, Sheep</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture

supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	BALB/c mouse mammary tumor cells
External Database Links	UniProt: P23229 Related reagents Entrez Gene: 3655 ITGA6 Related reagents
RRID	AB_321462
Fusion Partners	Spleen cells from immunized Sprague-Dawley rats were fused with cells of the SP2/0 mouse myeloma cell line
Specificity	<p>Rat anti Human CD49f antibody, clone NKI-GoH3 recognizes CD49f, also known as the VLA-6 alpha chain. CD49f is a 1107 amino acid ~120 kDa cell surface glycoprotein that forms distinct complexes with CD29 (VLA beta-chain), resulting in the VLA-6 (alpha-6 beta-1) complex, expressed on human platelets, or with the beta-4 integrin resulting in the alpha-6 beta-4 complex expressed on various human epithelial cells.</p> <p>Rat anti Human CD49f antibody, clone NKI-GoH3 reacts with platelets, megakaryocytes, T lymphocytes and common acute lymphoblastic leukemia cells (alpha-6 beta-1). In immunohistology the monoclonal antibody reacts with epithelial cells of a variety of tissues, peripheral nerves, microvascular endothelial cells, placenta cyto- and syncytiotrophoblasts. VLA-6 is an important mediator of cell binding to laminin.</p> <p>Rat anti Human CD49f antibody, clone NKI-GoH3 blocks the binding of cells to the E8 fragment of laminin (Sonnenberg <i>et al.</i> 1998).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ platelets in 100ul.
Histology Positive Control Tissue	Human Tonsil
References	<ol style="list-style-type: none">1. Jensen, K.B. <i>et al.</i> (2010) Assaying proliferation and differentiation capacity of stem cells using disaggregated adult mouse epidermis. Nat Protoc. 5 (5): 898-911.2. Soligo, D. <i>et al.</i> (1989) Immunohistochemical reactivity on bone marrow and tissues of

- anti-VLA antibodies in the platelet panel, in Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W. *et al.* Oxford University Press p1029-1032.
3. Sonnenberg, A. *et al.* (1986) Development of mouse mammary gland: identification of stages in differentiation of luminal and myoepithelial cells using monoclonal antibodies and polyvalent antiserum against keratin. [J Histochem Cytochem. 34 \(8\): 1037-46.](#)
 4. Sonnenberg, A. *et al.* (1987) A complex of platelet glycoproteins Ic and IIa identified by a rat monoclonal antibody. [J Biol Chem. 262 \(21\): 10376-83.](#)
 5. Hemler, M.E. *et al.* (1988) Multiple very late antigen (VLA) heterodimers on platelets. Evidence for distinct VLA-2, VLA-5 (fibronectin receptor), and VLA-6 structures. [J Biol Chem. 263 \(16\): 7660-5.](#)
 6. Galkowska, H. *et al.* (1996) Reactivity of antibodies directed against human antigens with surface markers on canine leukocytes. [Vet Immunol Immunopathol. 53 \(3-4\): 329-34.](#)
 7. Sonnenberg, A. *et al.* (1988) Laminin receptor on platelets is the integrin VLA-6. [Nature. 336 \(6198\): 487-9.](#)
 8. Sonnenberg, A. *et al.* (1990) Integrin recognition of different cell-binding fragments of laminin (P1, E3, E8) and evidence that alpha 6 beta 1 but not alpha 6 beta 4 functions as a major receptor for fragment E8. [J Cell Biol. 110 \(6\): 2145-55.](#)
 9. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)
 10. Sonnenberg, A. *et al.* (1990) The alpha 6 beta 1 (VLA-6) and alpha 6 beta 4 protein complexes: tissue distribution and biochemical properties. [J Cell Sci. 96 \(Pt 2\): 207-17.](#)
 11. Sonnenberg, A. *et al.* (1988) Identification and characterization of a novel antigen complex on mouse mammary tumor cells using a monoclonal antibody against platelet glycoprotein Ic. [J Biol Chem. 263 \(28\): 14030-8.](#)
 12. Le Bellego, F. *et al.* (2005) Cytoskeleton reorganization mediates alpha6beta1 integrin-associated actions of laminin on proliferation and survival, but not on steroidogenesis of ovine granulosa cells. [Reprod Biol Endocrinol. 3: 19.](#)
 13. Anderson, C. *et al.* (2009) Sonic hedgehog-dependent synthesis of laminin alpha1 controls basement membrane assembly in the myotome. [Development. 136: 3495-504.](#)
 14. Collins, C.A. *et al.* (2011) Reprogramming adult dermis to a neonatal state through epidermal activation of β -catenin [Development. 138: 5189-99.](#)
 15. Moreira, M. L. *et al.* (2016) Vaccination against canine leishmaniosis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. [Veterinary Parasitology. 15 Feb \[Epub ahead of print\]](#)
 16. Mastrogiannaki M *et al.* (2016) β -catenin stabilization in skin fibroblasts causes fibrotic lesions by preventing adipocyte differentiation of the reticular dermis. [J Invest Dermatol. pii: S0022-202X\(16\)00489-9. \[Epub ahead of print\]](#)
 17. Schäfer, G. *et al.* (2013) The role of inflammation in HPV infection of the Oesophagus. [BMC Cancer. 13: 185.](#)
 18. Peuhu, E. *et al.* (2017) Integrin beta 1 inhibition alleviates the chronic hyperproliferative dermatitis phenotype of SHARPIN-deficient mice [PLOS ONE. 12 \(10\): e0186628.](#)
 19. Rayagiri, S.S. *et al.* (2018) Basal lamina remodeling at the skeletal muscle stem cell niche mediates stem cell self-renewal. [Nat Commun. 9 \(1\): 1075.](#)
 20. Loureiro, J. *et al.* (2019) Conjugation of the T1 sequence from CCN1 to fibrin

hydrogels for therapeutic vascularization. [Mater Sci Eng C Mater Biol Appl. 104: 109847.](#)
21. Ikeda, A. *et al.* (2020) Follistatin expressed in mechanically-damaged salivary glands of male mice induces proliferation of CD49f⁺ cells. [Sci Rep. 10 \(1\): 19959.](#)

Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.
------------------------	--

Storage	Store at +4°C or at -20°C if preferred.
----------------	---

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA699 10040
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M368788:200529'

Printed on 18 Jan 2024