

Datasheet: MCA699T

Description:	RAT ANTI HUMAN CD49f
Specificity:	CD49f
Other names:	INTEGRIN ALPHA 6 CHAIN, VLA-6
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	NKI-GoH3
Isotype:	IgG2a
Quantity:	25 µg

Product Details

RRID AB_1102169

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Mouse, Dog, Pig, Cynomolgus monkey, Sheep
N.B. Antibody reactivity and working conditions may vary between species.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide

Carrier Free

Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	BALB/c mouse mammary tumor cells
External Database Links	<p>UniProt: P23229 Related reagents</p> <p>Entrez Gene: 3655 ITGA6 Related reagents</p>
Fusion Partners	Spleen cells from immunized Sprague-Dawley rats were fused with cells of the SP2/0 mouse myeloma cell line
Specificity	<p>Rat anti Human CD49f antibody, clone NKI-GoH3 recognizes CD49f, also known as the VLA-6 alpha chain. CD49f is a 1107 amino acid ~120 kDa cell surface glycoprotein that forms distinct complexes with CD29 (VLA beta-chain), resulting in the VLA-6 (alpha-6 beta-1) complex, expressed on human platelets, or with the beta-4 integrin resulting in the alpha-6 beta-4 complex expressed on various human epithelial cells.</p> <p>Rat anti Human CD49f antibody, clone NKI-GoH3 reacts with platelets, megakaryocytes, T lymphocytes and common acute lymphoblastic leukemia cells (alpha-6 beta-1). In immunohistology the monoclonal antibody reacts with epithelial cells of a variety of tissues, peripheral nerves, microvascular endothelial cells, placenta cyto- and syncytiotrophoblasts. VLA-6 is an important mediator of cell binding to laminin.</p> <p>Rat anti Human CD49f antibody, clone NKI-GoH3 blocks the binding of cells to the E8 fragment of laminin (Sonnenberg et al. 1998).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ platelets in 100ul.
Histology Positive Control Tissue	Human Tonsil
References	<ol style="list-style-type: none"> Jensen, K.B. <i>et al.</i> (2010) Assaying proliferation and differentiation capacity of stem cells using disaggregated adult mouse epidermis. Nat Protoc. 5 (5): 898-911. Soligo, D. <i>et al.</i> (1989) Immunohistochemical reactivity on bone marrow and tissues of anti-VLA antibodies in the platelet panel, in Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W. <i>et al.</i> Oxford University Press p1029-1032. Sonnenberg, A. <i>et al.</i> (1986) Development of mouse mammary gland: identification of stages in differentiation of luminal and myoepithelial cells using monoclonal antibodies and polyvalent antiserum against keratin. J Histochem Cytochem. 34 (8): 1037-46. Sonnenberg, A. <i>et al.</i> (1987) A complex of platelet glycoproteins Ic and IIa identified by a rat monoclonal antibody. J Biol Chem. 262 (21): 10376-83. Hemler, M.E. <i>et al.</i> (1988) Multiple very late antigen (VLA) heterodimers on platelets. Evidence for distinct VLA-2, VLA-5 (fibronectin receptor), and VLA-6 structures. J Biol Chem. 263 (16): 7660-5. Galkowska, H. <i>et al.</i> (1996) Reactivity of antibodies directed against human antigens with surface markers on canine leukocytes. Vet Immunol Immunopathol. 53 (3-4): 329-34. Sonnenberg, A. <i>et al.</i> (1988) Laminin receptor on platelets is the integrin VLA-6. Nature. 336 (6198): 487-9. Sonnenberg, A. <i>et al.</i> (1990) Integrin recognition of different cell-binding fragments of laminin

- (P1, E3, E8) and evidence that alpha 6 beta 1 but not alpha 6 beta 4 functions as a major receptor for fragment E8. [J Cell Biol. 110 \(6\): 2145-55.](#)
9. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)
10. Sonnenberg, A. *et al.* (1990) The alpha 6 beta 1 (VLA-6) and alpha 6 beta 4 protein complexes: tissue distribution and biochemical properties. [J Cell Sci. 96 \(Pt 2\): 207-17.](#)
11. Sonnenberg, A. *et al.* (1988) Identification and characterization of a novel antigen complex on mouse mammary tumor cells using a monoclonal antibody against platelet glycoprotein Ic. [J Biol Chem. 263 \(28\): 14030-8.](#)
12. Le Bellego, F. *et al.* (2005) Cytoskeleton reorganization mediates alpha6beta1 integrin-associated actions of laminin on proliferation and survival, but not on steroidogenesis of ovine granulosa cells. [Reprod Biol Endocrinol. 3: 19.](#)
13. Anderson, C. *et al.* (2009) Sonic hedgehog-dependent synthesis of laminin alpha1 controls basement membrane assembly in the myotome. [Development. 136: 3495-504.](#)
14. Collins, C.A. *et al.* (2011) Reprogramming adult dermis to a neonatal state through epidermal activation of β -catenin [Development. 138: 5189-99.](#)
15. Moreira, M. L. *et al.* (2016) Vaccination against canine leishmaniosis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. [Veterinary Parasitology. 15 Feb \[Epub ahead of print\]](#)
16. Mastrogiannaki M *et al.* (2016) β -catenin stabilization in skin fibroblasts causes fibrotic lesions by preventing adipocyte differentiation of the reticular dermis. [J Invest Dermatol. pii: S0022-202X\(16\)00489-9. \[Epub ahead of print\]](#)
17. Schäfer, G. *et al.* (2013) The role of inflammation in HPV infection of the Oesophagus. [BMC Cancer. 13: 185.](#)
18. Peuhu, E. *et al.* (2017) Integrin beta 1 inhibition alleviates the chronic hyperproliferative dermatitis phenotype of SHARPIN-deficient mice [PLOS ONE. 12 \(10\): e0186628.](#)
19. Rayagiri, S.S. *et al.* (2018) Basal lamina remodeling at the skeletal muscle stem cell niche mediates stem cell self-renewal. [Nat Commun. 9 \(1\): 1075.](#)

Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)

[FITC](#)

Goat Anti Rat IgG (STAR69...) [FITC](#)
Goat Anti Rat IgG (STAR131...) [Alk. Phos.](#), [Biotin](#)
Goat Anti Rat IgG (STAR73...) [RPE](#)
Rabbit Anti Rat IgG (STAR21...) [HRP](#)
Goat Anti Rat IgG (STAR72...) [HRP](#)
Rabbit Anti Rat IgG (STAR16...) [DyLight®800](#)
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight®649](#), [DyLight®800](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M353237:190411'

Printed on 11 Apr 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)