

Datasheet: MCA708 **BATCH NUMBER 147003**

Description:MOUSE ANTI HUMAN CD10Specificity:CD103Format:PurifiedProduct Type:Monoclonal AntibodyClone:2G5.1Isotype:IgG2a		
Format: Purified Product Type: Monoclonal Antibody Clone: 2G5.1	Description:	MOUSE ANTI HUMAN CD103
Product Type: Monoclonal Antibody Clone: 2G5.1	Specificity:	CD103
Clone: 2G5.1	Format:	Purified
	Product Type:	Monoclonal Antibody
Isotype: IgG2a	Clone:	2G5.1
	Isotype:	lgG2a
Quantity: 0.2 mg	Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.biorad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/50 - 1/100
Immunohistology - Frozen (1)	-			1/50 - 1/100
Immunohistology - Paraffin				
ELISA			•	
Immunoprecipitation	•			
Western Blotting			•	
Immunofluorescence	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross	Reacts with: Cynomolgus monkey, Rhesus Monkey
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross
	reactivity is derived from testing within our laboratories, peer-reviewed publications or
	personal communications from the originators. Please refer to references indicated for
	further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human intestinal lymphocytes
External Database Links	UniProt: P38570 Related reagents Entrez Gene: 3682 ITGAE Related reagents
RRID	AB_321984
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line
Specificity	Mouse anti Human CD103 antibody, clone 2G5.1 recognizes the human CD103 cell surface antigen, also known as the alpha ^E integrin, HML-1 antigen, Mucosal lymphocyte 1 antigen or Integrin alpha-E. CD103 is a 1161 amino acid, ~130 kDa single pass type 1 transmembrane glycoprotein.
	Mouse anti Human CD103 antibody, clone 2G5.1 reacts with various subsets of human intra-epithelial lymphocytes:- 40% of lamina propria T-cells; 30% of mesenteric lymphoblasts and some lymphocytes from other mucosa. It recognises less than 0.1% bone marrow cells and less than 0.7% PBL.
Flow Cytometry	Use 10ul of the suggested working dilution to label 5 x 10^5 cells in 100ul.
Histology Positive Control Tissue	Intestine
References	 Cerf-Bensussan, N. et al. (1987) A monoclonal antibody (HML-1) defining a novel membrane molecule present on human intestinal lymphocytes. <u>Eur J Immunol. 17 (9): 1279-85.</u> Cerf-Bensussan, N. et al. (1986) A monoclonal antibody specific for rat intestinal lymphocytes. <u>J Immunol. 136 (1): 76-82.</u> Sopper, S. et al. (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. <u>Cytometry. 29 (4): 351-62.</u> Yoshino, N. et al. (2000) Upgrading of flow cytometric analysis for absolute counts,

cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. Exp. Anim. 49: 97-110.

- 5. Swan, Z.D. *et al.* (2015) Macrophage accumulation in gut mucosa differentiates AIDS from chronic SIV infection in rhesus macaques. <u>Eur J Immunol. Nov 9. [Epub ahead of print]</u>
- 6. Presicce P *et al.* (2012) Myeloid dendritic cells isolated from tissues of SIV-infected Rhesus macaques promote the induction of regulatory T cells. <u>AIDS. 26 (3): 263-73.</u>
- 7. Kutleša, S. *et al.* (2002) E-cadherin-mediated interactions of thymic epithelial cells with CD103+ thymocytes lead to enhanced thymocyte cell proliferation. <u>J Cell Sci. 115 (Pt 23):</u> 4505-15.

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA708 10041		
Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) RPE

Rabbit Anti Mouse IgG (STAR13...) HRP

Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody sales uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M373004:200811'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint