

Datasheet: MCA708T

| Description:  | MOUSE ANTI HUMAN CD103 |
|---------------|------------------------|
| Specificity:  | CD103                  |
| Format:       | Purified               |
| Product Type: | Monoclonal Antibody    |
| Clone:        | 2G5.1                  |
| Isotype:      | lgG2a                  |
| Quantity:     | 25 μg                  |
|               |                        |

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                              | Yes | No | Not Determined | Suggested Dilution |
|------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry               | •   |    |                | 1/50 - 1/100       |
| Immunohistology - Frozen (1) | •   |    |                | 1/50 - 1/100       |
| Immunohistology - Paraffin   |     | •  |                |                    |
| ELISA                        |     |    | •              |                    |
| Immunoprecipitation          | •   |    |                |                    |
| Western Blotting             |     |    | •              |                    |
| Immunofluorescence           |     |    |                |                    |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

| Target Species              | Human                                                                                                                                                                                                                                                                                                                                      |
|-----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Species Cross<br>Reactivity | Reacts with: Cynomolgus monkey, Rhesus Monkey <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form                | Purified IgG - liquid                                                                                                                                                                                                                                                                                                                      |

| Preparation                          | Purified IgG prepared by affinity chromatography on Protein A from ascites                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|--------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Buffer Solution                      | Phosphate buffered saline                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Preservative<br>Stabilisers          | <0.1% Sodium Azide (NaN <sub>3</sub> ) 0.5% Bovine Serum Albumin                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Approx. Protein<br>Concentrations    | IgG concentration 1.0 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Immunogen                            | Human intestinal lymphocytes                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| External Database<br>Links           | UniProt: P38570 Related reagents  Entrez Gene: 3682 ITGAE Related reagents                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| RRID                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|                                      | AB_1100549                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Fusion Partners                      | Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Specificity                          | <b>Mouse anti Human CD103 antibody, clone 2G5.1</b> recognizes the human CD103 cell surface antigen, also known as the alpha <sup>E</sup> integrin, HML-1 antigen, Mucosal lymphocyte 1 antigen or Integrin alpha-E. CD103 is a 1161 amino acid, ~130 kDa single pass type 1 transmembrane glycoprotein.                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|                                      | Mouse anti Human CD103 antibody, clone 2G5.1 reacts with various subsets of human intra-epithelial lymphocytes:- 40% of lamina propria T-cells; 30% of mesenteric lymphoblasts and some lymphocytes from other mucosa. It recognises less than 0.1% bone marrow cells and less than 0.7% PBL.                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Flow Cytometry                       | Use 10ul of the suggested working dilution to label 5 x $10^5$ cells in 100ul.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Histology Positive<br>Control Tissue | Intestine                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| References                           | <ol> <li>Cerf-Bensussan, N. et al. (1987) A monoclonal antibody (HML-1) defining a novel membrane molecule present on human intestinal lymphocytes. Eur J Immunol. 17 (9): 1279-85.</li> <li>Cerf-Bensussan, N. et al. (1986) A monoclonal antibody specific for rat intestinal lymphocytes. J Immunol. 136 (1): 76-82.</li> <li>Sopper, S. et al. (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. Cytometry. 29 (4): 351-62.</li> <li>Yoshino, N. et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (Macaca fascicularis) by using anti-human cross-reactive antibodies. Exp. Anim. 49: 97-110.</li> </ol> |

- 5. Swan, Z.D. *et al.* (2015) Macrophage accumulation in gut mucosa differentiates AIDS from chronic SIV infection in rhesus macaques. <u>Eur J Immunol. Nov 9. [Epub ahead of print]</u>
- 6. Presicce P *et al.* (2012) Myeloid dendritic cells isolated from tissues of SIV-infected Rhesus macaques promote the induction of regulatory T cells. <u>AIDS. 26 (3): 263-73.</u>
- 7. Kutleša, S. *et al.* (2002) E-cadherin-mediated interactions of thymic epithelial cells with CD103+ thymocytes lead to enhanced thymocyte cell proliferation. <u>J Cell Sci. 115 (Pt 23):</u> 4505-15.
- 8. Nugent, J.L. *et al.* (2021) A nonhuman primate model of vertical sleeve gastrectomy facilitates mechanistic and translational research in human obesity. <u>iScience. 24 (12):</u> 103421.

#### **Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

| Guarantee                        | 12 months from date of despatch                                                                                               |
|----------------------------------|-------------------------------------------------------------------------------------------------------------------------------|
| Health And Safety<br>Information | Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf |
| Regulatory                       | For research purposes only                                                                                                    |

## Related Products

### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

Human Anti Mouse IgG2a (HCA037...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR76...)

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

**Recommended Negative Controls** 

### MOUSE IgG2a NEGATIVE CONTROL (MCA929)

Email: antibody\_sales\_us@bio-rad.com

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_uk@bio-rad.com

Page 3 of 4

Email: antibody\_sales\_de@bio-rad.com

## Printed on 13 May 2022

© 2022 Bio-Rad Laboratories Inc | Legal | Imprint