

Datasheet: MCA715PE

Description:	RAT ANTI HUMAN CD59:RPE
Specificity:	CD59
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	YTH53.1
Isotype:	lgG2b
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjuga	ated to R. Phycoerythrin	(RPE) - lyophilized
Reconstitution	Reconstitute with 1	ml distilled water	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
		.00	
reparation	Purified IgG prepare supernatant	ed by affinity chromatogi	raphy on Protein G f
		ed by affinity chromatog	raphy on Protein G f
Buffer Solution Preservative	supernatant	ed by affinity chromatogo	raphy on Protein G f
Preparation Buffer Solution Preservative Stabilisers	supernatant Phosphate buffered	ed by affinity chromatogod I saline	raphy on Protein G f

1			_	
lm	ΠIL	mo	u	en

Human Peripheral Blood T cells.

External Database

Links

UniProt:

P13987 Related reagents

Entrez Gene:

966 CD59 Related reagents

Synonyms

MIC11, MIN1, MIN2, MIN3, MSK21

RRID

AB 321513

Fusion Partners

Spleen cells from an immunised DA rat were fused with cells of the Y3/Ag1.2.3 rat myeloma cell line.

Specificity

Rat anti Human CD59 antibody, clone YTH53.1 recognizes the human CD59 cell surface antigen, also known as MAC-inhibitory protein, 1F5 antigen, 20 kDa homologous restriction factor, MEM43 antigen, membrane attack complex inhibition factor, membrane inhibitor of reactive lysis or protectin. CD59 is produced as a 103 amino acid proprotein, further processed to produce an 18-20 kDa GPI linked glycoprotein broadly expressed by human leucocytes and erythrocytes.

Mutation of the CD59 gene can result in the presentation of Hemolytic anemia, CD59-mediated, with or without polyneuropathy (<u>HACD59</u>) a disease characterized by infantile onset of a relapsing-remitting polyneuropathy manifest as limb muscle weakness, and hyporeflexia.

Rat anti Human CD59 antibody, clone YTH53.1 blocks the normal function of CD59 (Walsh *et al.* 1991).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. We do not recommend using this antibody in lysed whole blood techniques due to inhibition of staining caused by CD59 expression on erythrocytes.

References

- 1. Walsh, L.A. *et al.* (1991) Transfection of human CD59 complementary DNA into rat cells confers resistance to human complement. Eur J Immunol. 21 (3): 847-50.
- 2. Davies, A. *et al.* (1995) Identification of MIC 11 antigen as an epitope of the CD59 molecule. Immunology. 85 (2): 220-7.
- 3. Hosokawa, M. *et al.* (2004) Human oligodendroglial cells express low levels of C1 inhibitor and membrane cofactor protein mRNAs. <u>J Neuroinflammation</u>. 1: 17.
- 4. Ajona, D. *et al.* (2007) Down-regulation of human complement factor H sensitizes non-small cell lung cancer cells to complement attack and reduces *in vivo* tumor growth. J Immunol. 178 (9): 5991-8.
- 5. Ellison BS *et al.* (2007) Complement susceptibility in glutamine deprived breast cancer cells. Cell Div. 2: 20.
- 6. Koch, N. *et al.* (2009) IL-10 protects monocytes and macrophages from complement-mediated lysis. J Leukoc Biol. 86: 155-66.

7. Yang, P. et al. (2009) Expression and modulation of RPE cell membrane complement regulatory proteins. Invest Ophthalmol Vis Sci. 50: 3473-81.

8. Zhu, C. et al. (2020) Isatuximab Acts Through Fc-Dependent, Independent, and Direct Pathways to Kill Multiple Myeloma Cells. Front Immunol. 11: 1771.

Storage

This product is shipped at ambient temperature.

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA715PE 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

RAT IgG2b NEGATIVE CONTROL:RPE (MCA6006PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M441292:250523'

Printed on 24 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint