

Datasheet: MCA725G BATCH NUMBER 161369

| Description: | MOUSE ANTI HUMAN TALIN |
|---------------|------------------------|
| Specificity: | TALIN |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | TA205 |
| Isotype: | lgG1 |
| Quantity: | 0.2 mg |
| | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|---------------------------|
| Flow Cytometry (1) | • | | | 1/25 - 1/100 |
| Immunohistology - Frozen | | | | |
| Immunohistology - Paraffin | | | • | |
| ELISA | | | • | |
| Immunoprecipitation | | | | |
| Western Blotting | | | | |
| Immunofluorescence | | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm® (Product Code <u>BUF09</u>) for this purpose.

| Target Species | Human |
|-----------------------------|---|
| Species Cross Reactivity | Reacts with: Rabbit, Chicken N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form | Purified IgG - liquid |

| Preparation | Purified IgG prepared by affinity chromatography on Protein G | | | | |
|--------------------------------------|--|--|--|--|--|
| Buffer Solution | Phosphate buffered saline | | | | |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) | | | | |
| Carrier Free | Yes | | | | |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml | | | | |
| Immunogen | Human platelet talin. | | | | |
| External Database Links | UniProt: Q9Y490 Related reagents Q9Y4G6 Related reagents Entrez Gene: | | | | |
| | 7094 TLN1 Related reagents 83660 TLN2 Related reagents | | | | |
| Synonyms | KIAA0320, KIAA1027, TLN | | | | |
| RRID | AB_2233001 | | | | |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the SP2/0 mouse myeloma cell line. | | | | |
| Specificity | Mouse anti Human Talin antibody, clone TA205 recognizes human talin, a ~225 kDa cytoskeletal protein. Mouse anti Human Talin antibody, clone TA205 may be used to stain adhesion plaques in fixed and permeabilized fibroblasts. Mouse anti Human Talin antibody, clone recognizes an epitope between residues 139 and 433. | | | | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells | | | | |
| Histology Positive Control Tissue | Most tissues | | | | |
| Western Blotting | Clone TA205 has also been reported to work in Western blotting. | | | | |
| References | Porter, R.M. <i>et al.</i> (1993) Monoclonal antibodies to cytoskeletal proteins: an immunohistochemical investigation of human colon cancer. <u>J Pathol. 170 (4): 435-40.</u> Bolton, S.J. <i>et al.</i> (1997) Monoclonal antibodies recognizing the N- and C-terminal regions of talin disrupt actin stress fibers when microinjected into human fibroblasts. <u>Cell Motil Cytoskeleton. 36 (4): 363-76.</u> Zheng, P.P. <i>et al.</i> (2007) Hela I-CaD is implicated in the migration of endothelial cells/endothelial progenitor cells in human neoplasms. <u>Cell Adh Migr. 1: 84-91.</u> | | | | |

- 4. Vainionpää, N. *et al.* (2006) Laminin-10 and Lutheran blood group glycoproteins in adhesion of human endothelial cells. Am J Physiol Cell Physiol. 290: C764-75.
- 5. Filenius, S. (2003) Production of Fibronectin and Tenascin Isoforms and Their Role in the Adhesion of Human Immortalized Corneal Epithelial Cells <u>Invest Ophthalmol Vis Sci.</u> 44: 3317-25.
- 6. Calderwood, D.A. *et al.* (1999) The Talin head domain binds to integrin beta subunit cytoplasmic tails and regulates integrin activation. <u>J Biol Chem. 274: 28071-4.</u>
- 7. Cui, L. *et al.* (2009) c-Abl kinase is required for beta 2 integrin-mediated neutrophil adhesion. <u>J Immunol. 182: 3233-42.</u>
- 8. Koch, S. *et al.* (2013) Cathepsin D deficiency induces cytoskeletal changes and affects cell migration pathways in the brain. Neurobiol Dis. 50: 107-19.
- 9. Hasenson, S. *et al.* (2005) The immortalized human corneal epithelial cells adhere to laminin-10 by using Lutheran glycoproteins and integrin alpha3beta1. <u>Exp Eye Res. 81</u> (4): 415-21.
- 10. Braga, V.M. *et al.* (1995) Calcium-induced changes in distribution and solubility of cadherins, integrins and their associated cytoplasmic proteins in human keratinocytes. Cell Adhes Commun. 3 (3): 201-15.
- 11. Valmu, L. *et al.* (1999) The cytoskeletal association of CD11/CD18 leukocyte integrins in phorbol ester-activated cells correlates with CD18 phosphorylation. <u>Eur J Immunol. 29</u> (7): 2107-18.
- 12. Zhang C *et al.* (2013) c-Abl Kinase Is a Regulator of $\alpha\nu\beta$ 3 Integrin Mediated Melanoma A375 Cell Migration. <u>PLoS One. 8 (6): e66108.</u>

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA725G 10040 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...)

Goat Anti Mouse IgG (STAR70...)

Rabbit Anti Mouse IgG (STAR13...)

Goat Anti Mouse IgG (Fc) (STAR120...)

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381477:210512'

Printed on 17 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint