

Datasheet: MCA743F

BATCH NUMBER 1609

Description:	MOUSE ANTI HUMAN CD49b:FITC
Specificity:	CD49b
Other names:	INTEGRIN ALPHA 2 CHAIN, VLA-2
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	AK7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. The suggested working dilution is given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Cynomolgus monkey, Rhesus Monkey, Baboon
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
External Database Links	<p>UniProt: P17301 Related reagents</p> <p>Entrez Gene: 3673 ITGA2 Related reagents</p>
Synonyms	CD49B
RRID	AB_321447
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Human CD49b antibody, clone AK7 recognizes the integrin alpha 2 subunit, a ~160 kDa glycoprotein that non-covalently associates with the ~130 kDa integrin beta 1 subunit to form the VLA-2 complex.</p> <p>CD49b is expressed by platelets, long term cultivated T cells, approximately 50% of monocytes and most adherent cell lines.</p> <p>Mouse anti Human CD49b antibody, clone AK7 inhibits cell attachment to collagen.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
References	<ol style="list-style-type: none"> Favaloro, E.J. <i>et al.</i> (1990) Co-expression of haemopoietic antigens on vascular endothelial cells: a detailed phenotypic analysis. Br J Haematol. 74 (4): 385-94. Mazurov, A.V. <i>et al.</i> (1991) A monoclonal antibody, VM64, reacts with a 130 kDa glycoprotein common to platelets and endothelial cells: heterogeneity in antibody binding to human aortic endothelial cells. Thromb Haemost. 66 (4): 494-9. Yoshino N <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. Exp Anim. 49 (2): 97-110. Koutsogiannaki, S. & Kaloyianni, M. (2011) Effect of 17β-estradiol on adhesion of <i>Mytilus galloprovincialis</i> hemocytes to selected substrates. Role of alpha2 integrin subunit. Fish Shellfish Immunol. 31 (1): 73-80. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. Clin Exp Immunol. 127: 60-5.

6. Mörtberg A *et al.* (2016) Sensitive detection of platelet-specific antibodies with a modified MAIPA using biotinylated antibodies and streptavidin-coated beads. [J Immunol Methods. 434: 9-15.](#)
7. Rane, J.K. *et al.* (2016) Inhibition of the glucocorticoid receptor results in an enhanced miR-99a/100-mediated radiation response in stem-like cells from human prostate cancers. [Oncotarget. 7 \(32\): 51965-80.](#)
8. Hamaia, S.W. *et al.* (2012) Mapping of potent and specific binding motifs, GLOGEN and GVOGEA, for integrin $\alpha 1\beta 1$ using collagen toolkits II and III. [J Biol Chem. 287 \(31\): 26019-28.](#)
9. García-gareta, E. *et al.* (2019) Engineering the migration and attachment behaviour of primary dermal fibroblasts. [Biotechnol Bioeng. Jan 19 \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA743F>
10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M368865:200529'

Printed on 19 Jan 2024