

## Datasheet: MCA743T

**BATCH NUMBER 154563**

<b>Description:</b>	MOUSE ANTI HUMAN CD49b
<b>Specificity:</b>	CD49b
<b>Other names:</b>	INTEGRIN ALPHA 2 CHAIN, VLA-2
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AK7
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			1/500 - 1/1000
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

### Target Species

Human

### Species Cross Reactivity

Reacts with: Cynomolgus monkey, Rhesus Monkey, Baboon

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P17301</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3673</a> ITGA2    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CD49B
<b>RRID</b>	AB_2265115
<b>Fusion Partners</b>	Spleen cells from immunised mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD49b antibody, clone AK7</b> recognizes the integrin alpha 2 subunit, a ~160 kDa glycoprotein that non-covalently associates with the ~130 kDa integrin beta 1 subunit to form the VLA-2 complex.</p> <p>CD49b is expressed by platelets, long term cultivated T cells, approximately 50% of monocytes and most adherent cell lines.</p> <p>Mouse anti Human CD49b antibody, clone AK7 inhibits cell attachment to collagen.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>Histology Positive Control Tissue</b>	Tonsil
<b>References</b>	<ol style="list-style-type: none"> <li>Favaloro, E.J. <i>et al.</i> (1990) Co-expression of haemopoietic antigens on vascular endothelial cells: a detailed phenotypic analysis. <a href="#">Br J Haematol. 74 (4): 385-94.</a></li> <li>Mazurov, A.V. <i>et al.</i> (1991) A monoclonal antibody, VM64, reacts with a 130 kDa glycoprotein common to platelets and endothelial cells: heterogeneity in antibody binding to human aortic endothelial cells. <a href="#">Thromb Haemost. 66 (4): 494-9.</a></li> <li>Yoshino N <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts,</li> </ol>

cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)

4. Koutsogiannaki, S. & Kaloyianni, M. (2011) Effect of 17 $\beta$ -estradiol on adhesion of *Mytilus galloprovincialis* hemocytes to selected substrates. Role of alpha2 integrin subunit. [Fish Shellfish Immunol. 31 \(1\): 73-80.](#)

5. Cavers, M. *et al.* (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. [Clin Exp Immunol. 127: 60-5.](#)

6. Mörtberg A *et al.* (2016) Sensitive detection of platelet-specific antibodies with a modified MAIPA using biotinylated antibodies and streptavidin-coated beads. [J Immunol Methods. 434: 9-15.](#)

7. Rane, J.K. *et al.* (2016) Inhibition of the glucocorticoid receptor results in an enhanced miR-99a/100-mediated radiation response in stem-like cells from human prostate cancers. [Oncotarget. 7 \(32\): 51965-80.](#)

8. Hamaia, S.W. *et al.* (2012) Mapping of potent and specific binding motifs, GLOGEN and GVOGEA, for integrin  $\alpha$ 1 $\beta$ 1 using collagen toolkits II and III. [J Biol Chem. 287 \(31\): 26019-28.](#)

9. García-gareta, E. *et al.* (2019) Engineering the migration and attachment behaviour of primary dermal fibroblasts. [Biotechnol Bioeng. Jan 19 \[Epub ahead of print\].](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA743T>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),

Rabbit Anti Mouse IgG (STAR9...) [FITC](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...) [FITC](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

**Recommended Negative Controls**

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M368869:200529'

**Printed on 19 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)