

Datasheet: MCA743T

BATCH NUMBER 165377

| | |
|----------------------|-------------------------------|
| Description: | MOUSE ANTI HUMAN CD49b |
| Specificity: | CD49b |
| Other names: | INTEGRIN ALPHA 2 CHAIN, VLA-2 |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | AK7 |
| Isotype: | IgG1 |
| Quantity: | 25 µg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/50 - 1/100 |
| Immunohistology - Frozen (1) | ▪ | | | 1/500 - 1/1000 |
| Immunohistology - Paraffin | | ▪ | | |
| ELISA | | | ▪ | |
| Immunoprecipitation | | | ▪ | |
| Western Blotting | | | ▪ | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

| | |
|---------------------------------|--|
| Target Species | Human |
| Species Cross Reactivity | <p>Reacts with: Cynomolgus monkey, Rhesus Monkey, Baboon</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p> |

| | |
|--|--|
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |
| External Database Links | <p>UniProt: P17301 Related reagents</p> <p>Entrez Gene: 3673 ITGA2 Related reagents</p> |
| Synonyms | CD49B |
| RRID | AB_2265115 |
| Fusion Partners | Spleen cells from immunised mice were fused with cells of the mouse NS1 myeloma cell line. |
| Specificity | <p>Mouse anti Human CD49b antibody, clone AK7 recognizes the integrin alpha 2 subunit, a ~160 kDa glycoprotein that non-covalently associates with the ~130 kDa integrin beta 1 subunit to form the VLA-2 complex.</p> <p>CD49b is expressed by platelets, long term cultivated T cells, approximately 50% of monocytes and most adherent cell lines.</p> <p>Mouse anti Human CD49b antibody, clone AK7 inhibits cell attachment to collagen.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| Histology Positive Control Tissue | Tonsil |
| References | <ol style="list-style-type: none"> 1. Favaloro, E.J. <i>et al.</i> (1990) Co-expression of haemopoietic antigens on vascular endothelial cells: a detailed phenotypic analysis. Br J Haematol. 74 (4): 385-94. 2. Mazurov, A.V. <i>et al.</i> (1991) A monoclonal antibody, VM64, reacts with a 130 kDa glycoprotein common to platelets and endothelial cells: heterogeneity in antibody binding to human aortic endothelial cells. Thromb Haemost. 66 (4): 494-9. 3. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and |

L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. [Clin Exp Immunol. 127: 60-5.](#)

4. Koutsogiannaki, S. & Kaloyianni, M. (2011) Effect of 17 β -estradiol on adhesion of *Mytilus galloprovincialis* hemocytes to selected substrates. Role of alpha2 integrin subunit. [Fish Shellfish Immunol. 31 \(1\): 73-80.](#)

5. Hamaia, S.W. *et al.* (2012) Mapping of potent and specific binding motifs, GLOGEN and GVOGEA, for integrin $\alpha 1\beta 1$ using collagen toolkits II and III. [J Biol Chem. 287 \(31\): 26019-28.](#)

6. Mörtberg A *et al.* (2016) Sensitive detection of platelet-specific antibodies with a modified MAIPA using biotinylated antibodies and streptavidin-coated beads. [J Immunol Methods. 434: 9-15.](#)

7. Rane, J.K. *et al.* (2016) Inhibition of the glucocorticoid receptor results in an enhanced miR-99a/100-mediated radiation response in stem-like cells from human prostate cancers. [Oncotarget. 7 \(32\): 51965-80.](#)

8. García-gareta, E. *et al.* (2019) Engineering the migration and attachment behaviour of primary dermal fibroblasts. [Biotechnol Bioeng. Jan 19 \[Epub ahead of print\].](#)

| | |
|--------------------------------------|---|
| Storage | <p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p> |
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | <p>Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA743T</p> <p>10040</p> |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

| | |
|---|---|
| Goat Anti Mouse IgG (STAR77...) | HRP |
| Rabbit Anti Mouse IgG (STAR12...) | RPE |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | Alk. Phos. , HRP |
| Goat Anti Mouse IgG (STAR76...) | RPE |
| Goat Anti Mouse IgG (Fc) (STAR120...) | FITC , HRP |
| Rabbit Anti Mouse IgG (STAR13...) | HRP |
| Goat Anti Mouse IgG (STAR70...) | FITC |
| Goat Anti Mouse IgG (H/L) (STAR117...) | Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP |
| Rabbit Anti Mouse IgG (STAR9...) | FITC |

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M381489:210512'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)