

Datasheet: MCA74EL BATCH NUMBER 1704

Description:	RAT ANTI MOUSE CD11b:Low Endotoxin
Specificity:	CD11b
Other names:	INTEGRIN ALPHA M CHAIN, MAC-1
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	M1/70.15
Isotype:	IgG2b
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/100
Immunohistology - Frozen	•			
Immunohistology - Paraffin (1)	-			
ELISA			•	
Immunoprecipitation	•			
Western Blotting			•	
Immunofluorescence	-			
Functional Assays	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)PLP fixation is recommended for optimal results, see Whiteland et al. for details

Target Species	Mouse
Species Cross Reactivity	Reacts with: Human, Rabbit N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G from supernatant	n tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	
Carrier Free	Yes	
Endotoxin Level	< 0.01 EU/ug	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	T cell enriched splenocytes from B10 mice.	
External Database Links	UniProt: P05555 Related reagents Entrez Gene: 16409 Itgam Related reagents	
RRID	AB_2129277	
RRID Fusion Partners	AB_2129277 Spleen cells from immunized DA rats were fused with cells of the N myeloma cell line.	IS1/1.Ag4.1 mouse
	Spleen cells from immunized DA rats were fused with cells of the N	murine CD11b cell I, a differentiation e macrophages. Ind expression levels to express higher It to block iC3b binding It to as being suitable ested for use on formalin

- complement receptor. J Exp Med. 156 (4): 1000-9.
- 2. Whiteland, J.L. *et al.* (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. <u>J Histochem Cytochem. 43 (3): 313-20.</u>
- 3. Welt, F.G. *et al.* (2000) Neutrophil, not macrophage, infiltration precedes neointimal thickening in balloon-injured arteries. <u>Arterioscler Thromb Vasc Biol. 20 (12): 2553-8.</u>
- 4. Kapturczak, M.H. *et al* (2004) Heme oxygenase-1 modulates early inflammatory responses: evidence from the heme oxygenase-1-deficient mouse. <u>Am J Pathol. 165:</u> 1045-53.
- 5. Patel, P.C. and Harrison, R.E. (2008) Membrane ruffles capture C3bi-opsonized particles in activated macrophages. Mol Biol Cell. 19: 4628-39.
- 6. Huang, Q.Q. *et al* (2008) Role of H2-calponin in regulating macrophage motility and phagocytosis. <u>J Biol Chem. 283: 25887-99.</u>
- 7. Brochard, V. *et al* (2009) Infiltration of CD4+ lymphocytes into the brain contributes to neurodegeneration in a mouse model of Parkinson disease. J Clin Invest. 119: 182-92.
- 8. Hudcovic, T. *et al* (2009) Monocolonization with Bacteroides ovatus protects immunodeficient SCID mice from mortality in chronic intestinal inflammation caused by long-lasting dextran sodium sulfate treatment. <u>Physiol Res. 58: 101-10.</u>
- 9. Kroner, A. *et al* (2010) Ectopic T-cell specificity and absence of perforin and granzyme B alleviate neural damage in oligodendrocyte mutant mice. <u>Am J Pathol. 176: 549-55.</u>
- 10. Kanu, N. *et al.* (2010) The ATM cofactor ATMIN protects against oxidative stress and accumulation of DNA damage in the aging brain. <u>J Biol Chem. 285: 38534-42.</u>
- 11. Terrando, N. *et al.* (2010) The impact of IL-1 modulation on the development of lipopolysaccharide-induced cognitive dysfunction. Crit Care. 14 (3): R88.
- 12. Ferger, A.I. *et al* (2010) Effects of mitochondrial dysfunction on the immunological properties of microglia. <u>J Neuroinflammation</u>. 7: 45.
- 13. Gales, A. *et al* (2010) PPARgamma controls dectin-1 expression required for host antifungal defense against Candida albicans. <u>PLoS Pathog. 6: e1000714.</u>
- 14. Ghasemlou, N. *et al.* (2010) Mitogen-activated protein kinase-activated protein kinase 2 (MK2) contributes to secondary damage after spinal cord injury. <u>J Neurosci. 30:</u> 13750-9.
- 15. Lefevre, L. *et al.* (2010) PPARγ ligands switched high fat diet-induced macrophage M2b polarization toward M2a thereby improving intestinal Candida elimination. <u>PLoS One.</u> 5: e12828.
- 16. Samanta, J. *et al.* (2010) Noggin protects against ischemic brain injury in rodents. Stroke. 41: 357-62.
- 17. Yang, X. *et al* (2010) The role of the JAK2-STAT3 pathway in pro-inflammatory responses of EMF-stimulated N9 microglial cells. <u>J Neuroinflammation</u>. 7: 54.
- 18. Zhang, H. *et al.* (2010) A p38 mitogen-activated protein kinase-dependent mechanism of disinhibition in spinal synaptic transmission induced by tumor necrosis factor-alpha. <u>J Neurosci. 30: 12844-55.</u>
- 19. L'Episcopo, F. *et al.* (2010) Combining nitric oxide release with anti-inflammatory activity preserves nigrostriatal dopaminergic innervation and prevents motor impairment in a 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine model of Parkinson's disease. <u>J</u>
 Neuroinflammation. 7: 83.
- 20. Lebson, L. *et al.* (2010) Trafficking CD11b-positive blood cells deliver therapeutic genes to the brain of amyloid-depositing transgenic mice. <u>J Neurosci. 30: 9651-8.</u>

- 21. Kondo, Y. *et al.* (2011) Macrophages counteract demyelination in a mouse model of globoid cell leukodystrophy. J Neurosci. 31: 3610-24.
- 22. Geier, H. and Celli, J. (2011) Phagocytic Receptors Dictate Phagosomal Escape and Intracellular Proliferation of *Francisella tularensis*. <u>Infect Immun. 79: 2204-14.</u>
- 23. Redensek, A. *et al.* (2011) Expression and detrimental role of hematopoietic prostaglandin D synthase in spinal cord contusion injury. <u>Glia. 59: 603-14.</u>
- 24. Cunningham, K. *et al.* (2011) Conditional deletion of Ccm2 causes hemorrhage in the adult brain: a mouse model of human cerebral cavernous malformations. <u>Hum Mol Genet.</u> 20: 3198-206.
- 25. Fernández-Suárez,D. (2014) The monoacylglycerol lipase inhibitor JZL184 is neuroprotective and alters glial cell phenotype in the chronic MPTP mouse model Neurobiol Aging. 35: 2603-16.
- 26. Certo, M. *et al.* (2015) Activation of RXR/PPARγ underlies neuroprotection by bexarotene in ischemic stroke. Pharmacol Res. 102: 298-307.
- 27. Jones, R.S. *et al.* (2015) Inhibition of JAK2 attenuates the increase in inflammatory markers in microglia from APP/PS1 mice. Neurobiol Aging. 36 (10): 2716-24.
- 28. Nishikawa, K. *et al.* (2015) Resveratrol increases CD68⁺ Kupffer cells colocalized with adipose differentiation-related protein and ameliorates high-fat-diet-induced fatty liver in mice. Mol Nutr Food Res. 59 (6): 1155-70.
- 29. Khan, A.A. *et al.* (2015) Therapeutic Immunization with a Mixture of Herpes Simplex Virus Type 1 Glycoprotein D Derived "Asymptomatic" Human CD8+ T-cell Epitopes Decreases Spontaneous Ocular Shedding in Latently Infected HLA Transgenic Rabbits: Association with Low Frequency of Local PD-1+TIM-3+CD8+ Exhausted T Cells. <u>J Virol. pii: JVI.00788-15.</u>
- 30. Kim, B.W. *et al.* (2015) α-Asarone attenuates microglia-mediated neuroinflammation by inhibiting NF kappa B activation and mitigates MPTP-induced behavioral deficits in a mouse model of Parkinson's disease. Neuropharmacology. 97: 46-57.
- 31. Bains, M. & Roberts, J.L. (2016) Estrogen protects against dopamine neuron toxicity in primary mesencephalic cultures through an indirect P13K/Akt mediated astrocyte pathway. Neurosci Lett. 610: 79-85.
- 32. Amantea, D. *et al.* (2016) Azithromycin protects mice against ischemic stroke injury by promoting macrophage transition towards M2 phenotype. Exp Neurol. 275 Pt 1: 116-25.
- 33. Macrez, R. *et al.* (2016) Neuroendothelial NMDA receptors as therapeutic targets in experimental autoimmune encephalomyelitis. <u>Brain. 139 (Pt 9): 2406-19.</u>
- 34. Chen, Z.Z. *et al.* (2016) Memantine mediates neuroprotection via regulating neurovascular unit in a mouse model of focal cerebral ischemia. <u>Life Sci. 150: 8-14.</u>
- 35. Rich, M.C. *et al.* (2016) Site-targeted complement inhibition by a complement receptor 2-conjugated inhibitor (mTT30) ameliorates post-injury neuropathology in mouse brains. Neurosci Lett. 617: 188-94.
- 36. Werneburg, S. *et al.* (2016) Polysialylation and lipopolysaccharide-induced shedding of E-selectin ligand-1 and neuropilin-2 by microglia and THP-1 macrophages. <u>Glia. 64 (8): 1314-30.</u>
- 37. Amantea, D. *et al.* (2016) Neuroprotective Properties of a Macrolide Antibiotic in a Mouse Model of Middle Cerebral Artery Occlusion: Characterization of the Immunomodulatory Effects and Validation of the Efficacy of Intravenous Administration. Assay Drug Dev Technol. 14 (5): 298-307.
- 38. McCarthy, R.C. et al. (2016) Characterization of a novel adult murine immortalized

- microglial cell line and its activation by amyloid-beta. J Neuroinflammation. 13: 21.
- 39. Jiang, H. *et al.* (2017) Dense Intra-adipose Sympathetic Arborizations Are Essential for Cold-Induced Beiging of Mouse White Adipose Tissue. <u>Cell Metab. 26 (4): 686-692.e3.</u>
- 40. Zhang, J.C. *et al.* (2017) Prophylactic effects of sulforaphane on depression-like behavior and dendritic changes in mice after inflammation. <u>J Nutr Biochem. 39: 134-44.</u>
- 41. Petković, F. *et al.* (2017) Reduced cuprizone-induced cerebellar demyelination in mice with astrocyte-targeted production of IL-6 is associated with chronically activated, but less responsive microglia. J Neuroimmunol. 310: 97-102.
- 42. Martiskainen, H. *et al.* (2017) DHCR24 exerts neuroprotection upon inflammation-induced neuronal death. <u>J Neuroinflammation</u>. 14 (1): 215.
- 43. Olesen, M. N. *et al.* (2018) CD4 T cells react to local increase of α-synuclein in a pathology-associated variant-dependent manner and modify brain microglia in absence of brain pathology <u>Heliyon. 4 (1): e00513.</u>
- 44. Shin, D. *et al.* (2018) Bee Venom Phospholipase A2 Alleviate House Dust Mite-Induced Atopic Dermatitis-Like Skin Lesions by the CD206 Mannose Receptor. Toxins (Basel). 10 (4): 146.
- 45. Dos-Santos-Pereira, M. *et al.* (2018) Microglial glutamate release evoked by α-synuclein aggregates is prevented by dopamine. <u>Glia. 66 (11): 2353-2365.</u>
- 46. Price, B.R. *et al.* (2020) Therapeutic Trem2 activation ameliorates amyloid-beta deposition and improves cognition in the 5XFAD model of amyloid deposition. <u>J</u> Neuroinflammation. 17 (1): 238.
- 47. Lemaitre, D. *et al.* (2020) Collateral Sprouting of Peripheral Sensory Neurons Exhibits a Unique Transcriptomic Profile. Mol Neurobiol. 57 (10): 4232-49.
- 48. Lim, S.L. *et al.* (2020) Genetic Ablation of Hematopoietic Cell Kinase Accelerates Alzheimer's Disease-Like Neuropathology in Tg2576 Mice. <u>Mol Neurobiol. 57 (5):</u> 2447-60.
- 49. Paasila, P.J. *et al.* (2021) Ground state depletion microscopy as a tool for studying microglia-synapse interactions. <u>J Neurosci Res. 99 (6): 1515-32.</u>
- 50. Thiesler, H. *et al.* (2021) Polysialic acid and Siglec-E orchestrate negative feedback regulation of microglia activation. <u>Cell Mol Life Sci. 78 (4): 1637-53.</u>
- 51. Groh, J. *et al.* (2021) Immune modulation attenuates infantile neuronal ceroid lipofuscinosis in mice before and after disease onset. Brain Commun. 3 (2): fcab047.
- 52. Badr, M. *et al.* (2022) Expansion of regulatory T cells by CD28 superagonistic antibodies attenuates neurodegeneration in A53T-α-synuclein Parkinson's disease mice. <u>J</u> Neuroinflammation. 19 (1): 319.
- 53. Machado-Pereira, M. *et al.* (2022) Argonaute-2 protects the neurovascular unit from damage caused by systemic inflammation. <u>J Neuroinflammation</u>. <u>19 (1): 11.</u>
- 54. Karikari, A.A. *et al.* (2022) Neurodegeneration by α-synuclein-specific T cells in AAV-A53T-α-synuclein Parkinson's disease mice. <u>Brain Behav Immun. 101: 194-210.</u>
- 55. Zhao, Y. *et al.* (2022) Using Extracellular Vesicles Released by GDNF-Transfected Macrophages for Therapy of Parkinson Disease. <u>Cells. 11 (12): 1933.</u>
- 56. Ji, N. *et al.* (2022) VSIG4 Attenuates NLRP3 and Ameliorates Neuroinflammation via JAK2-STAT3-A20 Pathway after Intracerebral Hemorrhage in Mice. <u>Neurotox Res. 40 (1):</u> 78-88.
- 57. Jang, M. *et al.* (2023) Micrandilactone C, a Nortriterpenoid Isolated from Roots of Schisandra chinensis, Ameliorates Huntington's Disease by Inhibiting Microglial STAT3 Pathways <u>Cells. 12 (5): 786.</u>

- 58. Tseng, K.Y. *et al.* (2023) Augmenting hematoma-scavenging capacity of innate immune cells by CDNF reduces brain injury and promotes functional recovery after intracerebral hemorrhage. Cell Death Dis. 14 (2): 128.
- 59. Wilton, D.K. *et al.* (2023) Microglia and complement mediate early corticostriatal synapse loss and cognitive dysfunction in Huntington's disease. <u>Nat Med. Oct 09 [Epub ahead of print].</u>
- 60. McFleder, R.L. *et al.* (2023) Brain-to-gut trafficking of alpha-synuclein by CD11c(+) cells in a mouse model of Parkinson's disease. Nat Commun. 14 (1): 7529.
- 61. Wang, Q. *et al.* (2018) Sarm1/Myd88-5 Regulates Neuronal Intrinsic Immune Response to Traumatic Axonal Injuries. <u>Cell Rep. 23 (3): 716-24.</u>
- 62. Türk Börü, Ü. *et al.* (2024) Alterations in the spinal cord, trigeminal nerve ganglion, and infraorbital nerve through inducing compression of the dorsal horn region at the upper cervical cord in trigeminal neuralgia. Brain Res. 1832: 148842.

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA74EL 10162
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...) DyLight®800

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR72...)

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

RPE

Rabbit Anti Rat IgG (STAR21...)

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®550, DyLight®650, DyLight®800

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

North & South Tel: +1 800 265 7376 Worldv

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

 Email: antibody_sales_de@bio-rad.com

Printed on 18 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint