

Datasheet: MCA752F

Description:	MOUSE ANTI GUINEA PIG CD8:FITC
Specificity:	CD8
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	CT6
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Guinea Pig						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin						
Approx. Protein	IgG concentration 0.1 mg/ml						

Concentrations

Immunogen Guinea pig peritoneal T-cells.

RRID AB_321399

Fusion Partners Spleen cells from immunized BALB/c mice were fused with cells of the X63.Ag8.653 mouse myeloma cell line.

Specificity **Mouse anti Guinea Pig CD8 antibody, clone CT6** reacts with guinea pig CD8 present on cytotoxic T-cells. CD8 comprises 2 subunits, alpha and beta and exists as either an alpha/alpha homodimer or an alpha/beta heterodimer. Sequence analysis suggests that guinea pig CD8 is more closely related to human than rat or mouse CD8 ([Nagarajan *et al.* 2004](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul

References

1. Tan, B.T.G. *et al.* (1985) Production of monoclonal antibodies defining guinea pig T-cell surface markers and a strain 13 Ia-like antigen: the value of immunohistological screening. [Hybridoma 4: 115-124.](#)
2. Steerenberg, P.A. *et al.* (1991) Tumour rejection after transfer of line 10 immunity is mediated by two T-cell populations. [Cancer Immunol. Immunother. 34: 103-110.](#)
3. Steerenberg, P.A. *et al.* (1990) Tumour infiltrating leucocytes (tils) during progressive tumour growth and BCG - mediated tumour regression. [Virchows Archiv Cell Pathol. 59: 185-194.](#)
4. Baker, D. *et al.* (1987) Changes in lymphocyte subsets after treatment with cyclophosphamide and during the development of contact sensitivity in the guinea pig. [Int J Immunopharmacol. 9 \(2\): 175-83.](#)
5. Antoniou, A.V. *et al.* (1986) Immunocytochemical identification and quantitation of mononuclear cells in the meninges during the development of chronic relapsing experimental allergic encephalomyelitis (CREAE) in the guinea pig. [Cell Immunol. 97 \(2\): 386-96.](#)
6. Liversidge, J. *et al.* (1987) EAU in the guinea pig: inhibition of cell-mediated immunity and Ia antigen expression by cyclosporin A. [Clin Exp Immunol. 69 \(3\): 591-600.](#)
7. Liversidge, J. & Forrester, J.V. (1988) Experimental autoimmune uveitis (EAU): immunophenotypic analysis of inflammatory cells in chorio retinal lesions. [Curr Eye Res. 7 \(12\): 1231-41.](#)
8. Debout, C. *et al.* (1991) The Kurloff cell in estrogenized guinea pigs as a CT7+ 8BE6-CT6- MR-1- CT10- IgM- lymphocyte with natural killer activity. [Nat Immun Cell Growth Regul. 10 \(6\): 327-35.](#)
9. Shang S *et al.* (2011) Activities of TMC207, rifampin, and pyrazinamide against Mycobacterium tuberculosis infection in guinea pigs. [Antimicrob Agents Chemother. 55 \(1\): 124-31.](#)
10. Lacy HM *et al.* (2011) Essential role for neutrophils in pathogenesis and adaptive immunity in *Chlamydia caviae* ocular infections. [Infect Immun. 79 \(5\): 1889-97.](#)
11. Komori, T. *et al.* (2011) A Microbial Glycolipid Functions as a New Class of Target Antigen for Delayed-type Hypersensitivity. [J Biol Chem. 286: 16800-6.](#)
12. Hiromatsu, K. *et al.* (2002) Induction of CD1-restricted immune responses in guinea

- pigs by immunization with mycobacterial lipid antigens. [J Immunol. 169: 330-9.](#)
13. Kramp, J.C. *et al.* (2011) The *in vivo* immunomodulatory effect of recombinant tumour necrosis factor-alpha in guinea pigs vaccinated with *Mycobacterium bovis* bacille Calmette-Guérin. [Clin Exp Immunol. 165: 110-20.](#)
14. Mishra, N.C. *et al.* (2010) Sulfur mustard induces immune sensitization in hairless guinea pigs. [Int Immunopharmacol. 10: 193-9.](#)
15. Schleiss, M.R. *et al.* (2007) Preconceptual administration of an alphavirus replicon UL83 (pp65 homolog) vaccine induces humoral and cellular immunity and improves pregnancy outcome in the guinea pig model of congenital cytomegalovirus infection. [J Infect Dis. 195: 789-98.](#)
16. Turner, O.C. *et al.* (2003) Immunopathogenesis of pulmonary granulomas in the guinea pig after infection with *Mycobacterium tuberculosis*. [Infect Immun. 71: 864-71.](#)
17. Wang, Y. *et al.* (2010) Local host response to chlamydial urethral infection in male guinea pigs. [Infect Immun. 78: 1670-81.](#)
18. Yang, H. *et al.* (2011) Three protein cocktails mediate delayed-type hypersensitivity responses indistinguishable from that elicited by purified protein derivative in the guinea pig model of *Mycobacterium tuberculosis* infection. [Infect Immun. 79: 716-23.](#)
19. Baldwin SL *et al.* (2012) The importance of adjuvant formulation in the development of a tuberculosis vaccine. [J Immunol. 188 \(5\): 2189-97.](#)
20. Obregón-Henao, A. Shang, s. Shanley, C.A. Basaraba, R.J. Caraway, M.L. Duncan, C.G. Ordway, D.J. Orme, I.M. (2013) Cortisone-Forced Reactivation of Weakly Acid Fast Positive *Mycobacterium Tuberculosis* in Guinea Pigs Previously Treated With Chemotherapy [Mycobacterial Diseases. 2: 116.](#)
21. Xia, J. *et al.* (2014) Virus-specific immune memory at peripheral sites of herpes simplex virus type 2 (HSV-2) infection in guinea pigs. [PLoS One. 9 \(12\): e114652.](#)
22. Jeevan A *et al.* (2013) Guinea pig skin, a model for epidermal cellular and molecular changes induced by UVR in vivo and in vitro: effects on *Mycobacterium bovis* Bacillus Calmette-Guérin vaccination. [Photochem Photobiol. 89 \(1\): 189-98.](#)
23. Gupta A *et al.* (2012) Protective efficacy of *Mycobacterium indicus pranii* against tuberculosis and underlying local lung immune responses in guinea pig model. [Vaccine. 30 \(43\): 6198-209.](#)
24. Gupta, A. *et al.* (2012) Efficacy of *Mycobacterium indicus pranii* immunotherapy as an adjunct to chemotherapy for tuberculosis and underlying immune responses in the lung. [PLoS One. 7 \(7\): e39215.](#)
25. Wu, W.H. *et al.* (2012) Immune status and the development of *Listeria monocytogenes* infection in aged and young guinea pigs. [Clin Invest Med. 35 \(5\): E309.](#)
26. Eckhardt, E. *et al.* (2023) Phosphatidylinositolmannoside vaccination induces lipid-specific Th1-responses and partially protects guinea pigs from *Mycobacterium tuberculosis* challenge. [Sci Rep. 13 \(1\): 18613.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA752F>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M410542:221028'

Printed on 23 Nov 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)