

Datasheet: MCA755SBV760

| Description: | MOUSE ANTI HUMAN CD27:StarBright Violet 760 | | |
|----------------------|---|--|--|
| Specificity: | CD27 | | |
| Format: | StarBright Violet 760 | | |
| Product Type: | Monoclonal Antibody | | |
| Clone: | LT27 | | |
| Isotype: | lgG2a | | |
| Quantity: | 100 TESTS/0.5ml | | |
| | | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | - | | | Neat |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Human |
|----------------|---|
| Species Cross | Reacts with: Cynomolgus monkey, Rhesus Monkey, Sheep |
| Reactivity | N.B. Antihody reactivity and working conditions may vary be |

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

| Product Form | Purified IgG conjugated to StarBright Violet 760 - liquid | | | | |
|-----------------|---|--|-------------------|--|--|
| | | | | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) | | |
| | StarBright Violet 760 | 403 | 754 | | |
| Preparation | Purified IgG prepare | Purified IgG prepared by ion exchange chromatography from ascite | | | |
| Buffer Solution | Phosphate buffered | saline | | | |

Preservative Stabilisers

0.09% Sodium Azide (NaN₃)

1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20 Approx. Protein For information on the concentration of our StarBright Dye conjugated reagents please Concentrations visit our FAQ page. **Immunogen** Normal human blood lymphocytes. **External Database UniProt:** Links P26842 Related reagents **Entrez Gene:** 939 CD27 Related reagents **Synonyms** TNFRSF7 **Fusion Partners** Spleen cells from immunized mice were fused with cells of the P3. X 63-Ag8-653 mouse myeloma cell line. **Specificity** Mouse anti human CD27, clone LT27, recognizes the human T-cell activation antigen, CD27. This antigen is expressed on activated T lymphocytes and belongs to a family of proteins associated with the nerve growth factor receptor. CD27 is expressed by thymocytes and peripheral T cells and also by EBV transformed B cell lines. Mouse anti human CD27, clone LT27 precipitates a ~55 kDa protein under reducing conditions or a ~120 kDa protein under non-reducing conditions. Flow Cytometry Use 5ul of the suggested working dilution to label 10⁶ cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application. References 1. Ravoet, A.M. et al. (1989) CD25 mAb: epitopes recognised, effect on lymphocyte activation, mediation of ADCC in Leucocyte Typing IV: White Cell Differentiation Antigens Eds Knapp, W. et al. Oxford University Press pp 408-411. 2. Boxall, S. et al. (2004) Abnormal cell surface antigen expression in individuals with variant CD45 splicing and histiocytosis. Pediatr Res. 55 (3): 478-84. 3. Griebel, P.J. et al. (2007) Cross-reactivity of mAbs to human CD antigens with sheep leukocytes. Vet Immunol Immunopathol. 119: 115-22. 4. Becker, P.D. et al. (2010) Generation of Human Antigen-Specific Monoclonal IgM Antibodies Using Vaccinated "Human Immune System" Mice PLoS One. 5(10). pii: e13137. 5. van Zelm, M.C. et al. (2007) Replication history of B lymphocytes reveals homeostatic proliferation and extensive antigen-induced B cell expansion. J Exp Med. 204: 645-55. 6. Steger, B. et al. (2014) CD4(+)and CD8(+)T-cell reactions against leukemia-associatedor minor-histocompatibility-antigens in AML-patients after allogeneic SCT. Immunobiology.

219 (4): 247-60.

| Storage | Store at +4°C. DO NOT FREEZE. |
|-------------------|---|
| | This product should be stored undiluted. |
| Guarantee | 12 months from date of despatch |
| Acknowledgements | This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts |
| Health And Safety | Material Safety Datasheet documentation #20471 available at: |
| Information | https://www.bio-rad-antibodies.com/SDS/MCA755SBV760 |
| | 20471 |
| Regulatory | For research purposes only |

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M437489:250314'

Printed on 10 Apr 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint