

Datasheet: MCA755SBY605

### **BATCH NUMBER 100007408**

Description:	MOUSE ANTI HUMAN CD27:StarBright Yellow 605
Specificity:	CD27
Format:	StarBright Yellow 605
Product Type:	Monoclonal Antibody
Clone:	LT27
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Human			
Reacts with: Cynomo	lgus monkey, Rhesus	Monkey, Sheep	
<b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publication personal communications from the originators. Please refer to references indicate further information.			
Purified IgG conjugate	ed to StarBright Yellow	<i>i</i> 605 - liquid	
Fluorophore	Excitation Max (nm)	Emission Max (nm)	
StarBright Yellow 605	572	606	
Purified IgG prepared	by affinity chromatog	raphy on DEAE from ascites	
Phosphate buffered s	aline		
	Reacts with: Cynomo  N.B. Antibody reactiv reactivity is derived fr personal communicat further information.  Purified IgG conjugate  Fluorophore StarBright Yellow 605  Purified IgG prepared	Reacts with: Cynomolgus monkey, Rhesus  N.B. Antibody reactivity and working conditi reactivity is derived from testing within our la personal communications from the originate further information.  Purified IgG conjugated to StarBright Yellow  Fluorophore Excitation Max (nm)  StarBright Yellow 605 572	Reacts with: Cynomolgus monkey, Rhesus Monkey, Sheep  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publication personal communications from the originators. Please refer to references indicate further information.  Purified IgG conjugated to StarBright Yellow 605 - liquid  Fluorophore Excitation Max (nm) Emission Max (nm)  StarBright Yellow 605 572 606  Purified IgG prepared by affinity chromatography on DEAE from ascites

Preservative Stabilisers 0.09% Sodium Azide (NaN<sub>3</sub>)1% Bovine Serum Albumin

0.1% Pluronic F680.1% PEG 33500.05% Tween 20

Immunogen

Normal human blood lymphocytes.

External Database Links

UniProt:

P26842 Related reagents

**Entrez Gene:** 

939 CD27 Related reagents

**Synonyms** 

TNFRSF7

**Fusion Partners** 

Spleen cells from immunized mice were fused with cells of the P3. X 63-Ag8-653 mouse myeloma cell line.

**Specificity** 

**Mouse anti human CD27, clone LT27,** recognizes the human T-cell activation antigen, CD27. This antigen is expressed on activated T lymphocytes and belongs to a family of proteins associated with the nerve growth factor receptor. CD27 is expressed by thymocytes and peripheral T cells and also by EBV transformed B cell lines.

Mouse anti human CD27, clone LT27 precipitates a ~55 kDa protein under reducing conditions or a ~120 kDa protein under non-reducing conditions.

**Flow Cytometry** 

Use 5ul of the suggested working dilution to label  $10^6$  cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Ravoet, A.M. *et al.* (1989) CD25 mAb: epitopes recognised, effect on lymphocyte activation, mediation of ADCC in Leucocyte Typing IV: White Cell Differentiation Antigens Eds Knapp, W. *et al.* Oxford University Press pp 408-411.
- 2. Boxall, S. *et al.* (2004) Abnormal cell surface antigen expression in individuals with variant CD45 splicing and histiocytosis. Pediatr Res. 55 (3): 478-84.
- 3. Griebel, P.J. *et al.* (2007) Cross-reactivity of mAbs to human CD antigens with sheep leukocytes. Vet Immunol Immunopathol. 119: 115-22.
- 4. Becker, P.D. *et al.* (2010) Generation of Human Antigen-Specific Monoclonal IgM Antibodies Using Vaccinated "Human Immune System" Mice <u>PLoS One. 5(10). pii:</u> <u>e13137.</u>
- 5. van Zelm, M.C. *et al.* (2007) Replication history of B lymphocytes reveals homeostatic proliferation and extensive antigen-induced B cell expansion. <u>J Exp Med. 204: 645-55.</u>
- 6. Steger, B. *et al.* (2014) CD4(+)and CD8(+)T-cell reactions against leukemia-associated-or minor-histocompatibility-antigens in AML-patients after allogeneic SCT. <u>Immunobiology.</u> 219 (4): 247-60.

**Storage** 

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted.

Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA755SBY605">https://www.bio-rad-antibodies.com/SDS/MCA755SBY605</a> 20471
Regulatory	For research purposes only

# **Related Products**

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

**Europe** Tel: +49 (0) 89 8090 95 21

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M415876:230116'

### Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint