

Datasheet: MCA756A647

**BATCH NUMBER 1608**

<b>Description:</b>	MOUSE ANTI HUMAN CD64:Alexa Fluor® 647
<b>Specificity:</b>	CD64
<b>Other names:</b>	FcRI
<b>Format:</b>	ALEXA FLUOR® 647
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	10.1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Baboon, Cynomolgus monkey, Rhesus Monkey  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Alexa Fluor® 647 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665

### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.05 mg/ml
<b>Immunogen</b>	Human monocytes
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P12314</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">2209</a>    FCGR1A    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	FCG1, FCGR1, IGFR1
<b>RRID</b>	AB_324802
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0-Ag14 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human CD64 antibody, clone 10.1</b> recognizes the human CD64 cell surface antigen, a ~75 kDa glycoprotein expressed by monocytes. The antigen acts as a high affinity receptor for human IgG, and is also known as FcRI.</p> <p>Mouse anti Human CD64 antibody, clone 10.1 blocks binding of immunoglobulin to FcRI.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Dougherty, G.J. <i>et al.</i> (1987) The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, 10.1. <a href="#">Eur J Immunol. 17 (10): 1453-9.</a></li> <li>2. Beekman, J.M. <i>et al.</i> (2004) Direct interaction between FcγRI (CD64) and periplakin controls receptor endocytosis and ligand binding capacity. <a href="#">Proc Natl Acad Sci U S A. 101: 10392-7.</a></li> <li>3. Tanaka, M. <i>et al.</i> (2009) Activation of Fc gamma RI on monocytes triggers differentiation into immature dendritic cells that induce autoreactive T cell responses. <a href="#">J Immunol. 183: 2349-55.</a></li> <li>4. Fet, N.G. <i>et al.</i> (2012) Reduction of activated macrophages after ischaemia-reperfusion injury diminishes oxidative stress and ameliorates renal damage. <a href="#">Nephrol Dial Transplant. 27 (8): 3149-55.</a></li> <li>5. Wagner, C. <i>et al.</i> (2008) T lymphocytes in acute bacterial infection: increased prevalence of CD11b(+) cells in the peripheral blood and recruitment to the infected site. <a href="#">Immunology. 125: 503-9.</a></li> <li>6. Eisenhardt, S.U. <i>et al.</i> (2009) Dissociation of pentameric to monomeric C-reactive protein on activated platelets localizes inflammation to atherosclerotic plaques. <a href="#">Circ Res.</a></li> </ol>

[105: 128-37.](#)

7. Fadlon, E. *et al.* (1998) Blood polymorphonuclear leukocytes from the majority of sickle cell patients in the crisis phase of the disease show enhanced adhesion to vascular endothelium and increased expression of CD64. [Blood. 91: 266-74.](#)

8. Scheinecker, C. *et al.* (1998) Initiation of the autologous mixed lymphocyte reaction requires the expression of costimulatory molecules B7-1 and B7-2 on human peripheral blood dendritic cells. [J Immunol. 161: 3966-73.](#)

9. Navarro-López, F. *et al.* (2003) Late T-lymphocyte and monocyte activation in coronary restenosis. Evidence for a persistent inflammatory/immune mechanism? [Rev Esp Cardiol. 56: 465-72.](#)

10. Liu M *et al.* (2011) Vitellogenin mediates phagocytosis through interaction with FcγR. [Mol Immunol. 49 \(1-2\): 211-8.](#)

11. Kapelski S *et al.* (2014) Assessment of the neutrophilic antibody-dependent respiratory burst (ADRB) response to *Plasmodium falciparum*. [J Leukoc Biol. 96 \(6\): 1131-42.](#)

12. Loi, A.L.T. *et al.* (2017) Proteomic profiling of peripheral blood neutrophils identifies two inflammatory phenotypes in stable COPD patients. [Respir Res. 18 \(1\): 100.](#)

13. Hristodorov, D. *et al.* (2016) Fully human MAP-fusion protein selectively targets and eliminates proliferating CD64(+) M1 macrophages. [Immunol Cell Biol. 94 \(5\): 470-8.](#)

14. Kahn, F. *et al.* (2008) Antibodies against a surface protein of *Streptococcus pyogenes* promote a pathological inflammatory response. [PLoS Pathog. 4 \(9\): e1000149.](#)

---

**Further Reading**

1. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Acknowledgements**

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information**      Material Safety Datasheet documentation #10041 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA756A647>  
10041

---

**Regulatory**                      For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South**      Tel: +1 800 265 7376

**America**              Fax: +1 919 878 3751

                            Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

                            Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

                            Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M368913:200529'

**Printed on 21 Feb 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)