

# Datasheet: MCA756F

**BATCH NUMBER 151134**

<b>Description:</b>	MOUSE ANTI HUMAN CD64:FITC
<b>Specificity:</b>	CD64
<b>Other names:</b>	FcRI
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	10.1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Baboon, Cynomolgus monkey, Rhesus Monkey <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Human monocytes
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P12314</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">2209</a>    FCGR1A    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	FCG1, FCGR1, IGFR1
<b>RRID</b>	AB_321799
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0-Ag14 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human CD64 antibody, clone 10.1</b> recognizes the human CD64 cell surface antigen, a ~75 kDa glycoprotein expressed by monocytes. The antigen acts as a high affinity receptor for human IgG, and is also known as FcRI.</p> <p>Mouse anti Human CD64 antibody, clone 10.1 blocks binding of immunoglobulin to FcRI.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Dougherty, G.J. <i>et al.</i> (1987) The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, 10.1. <a href="#">Eur J Immunol. 17 (10): 1453-9.</a></li> <li>2. Beekman, J.M. <i>et al.</i> (2004) Direct interaction between FcγRI (CD64) and periplakin controls receptor endocytosis and ligand binding capacity. <a href="#">Proc Natl Acad Sci U S A. 101: 10392-7.</a></li> <li>3. Tanaka, M. <i>et al.</i> (2009) Activation of Fc gamma RI on monocytes triggers differentiation into immature dendritic cells that induce autoreactive T cell responses. <a href="#">J Immunol. 183: 2349-55.</a></li> <li>4. Fet, N.G. <i>et al.</i> (2012) Reduction of activated macrophages after ischaemia-reperfusion injury diminishes oxidative stress and ameliorates renal damage. <a href="#">Nephrol Dial Transplant. 27 (8): 3149-55.</a></li> <li>5. Wagner, C. <i>et al.</i> (2008) T lymphocytes in acute bacterial infection: increased prevalence of CD11b(+) cells in the peripheral blood and recruitment to the infected site. <a href="#">Immunology. 125: 503-9.</a></li> <li>6. Eisenhardt, S.U. <i>et al.</i> (2009) Dissociation of pentameric to monomeric C-reactive protein on activated platelets localizes inflammation to atherosclerotic plaques. <a href="#">Circ Res.</a></li> </ol>

[105: 128-37.](#)

7. Fadlon, E. *et al.* (1998) Blood polymorphonuclear leukocytes from the majority of sickle cell patients in the crisis phase of the disease show enhanced adhesion to vascular endothelium and increased expression of CD64. [Blood. 91: 266-74.](#)
8. Scheinecker, C. *et al.* (1998) Initiation of the autologous mixed lymphocyte reaction requires the expression of costimulatory molecules B7-1 and B7-2 on human peripheral blood dendritic cells. [J Immunol. 161: 3966-73.](#)
9. Navarro-López, F. *et al.* (2003) Late T-lymphocyte and monocyte activation in coronary restenosis. Evidence for a persistent inflammatory/immune mechanism? [Rev Esp Cardiol. 56: 465-72.](#)
10. Liu M *et al.* (2011) Vitellogenin mediates phagocytosis through interaction with FcγR. [Mol Immunol. 49 \(1-2\): 211-8.](#)
11. Kapelski S *et al.* (2014) Assessment of the neutrophilic antibody-dependent respiratory burst (ADRB) response to *Plasmodium falciparum*. [J Leukoc Biol. 96 \(6\): 1131-42.](#)
12. Loi, A.L.T. *et al.* (2017) Proteomic profiling of peripheral blood neutrophils identifies two inflammatory phenotypes in stable COPD patients. [Respir Res. 18 \(1\): 100.](#)
13. Hristodorov, D. *et al.* (2016) Fully human MAP-fusion protein selectively targets and eliminates proliferating CD64(+) M1 macrophages. [Immunol Cell Biol. 94 \(5\): 470-8.](#)
14. Kahn, F. *et al.* (2008) Antibodies against a surface protein of *Streptococcus pyogenes* promote a pathological inflammatory response. [PLoS Pathog. 4 \(9\): e1000149.](#)

---

**Further Reading**

1. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10041 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA756F>  
10041

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M368916:200529'

**Printed on 29 Apr 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)