

Datasheet: MCA756PB

## **BATCH NUMBER 1807**

Description:	MOUSE ANTI HUMAN CD64:Pacific Blue®		
Specificity:	CD64		
Other names:	FcRI		
Format:	Pacific Blue®		
Product Type:	Monoclonal Antibody		
Clone:	10.1		
Isotype:	IgG1		
Quantity:	100 TESTS/1ml		

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Species Cross</b>
Reactivity

**Target Species** 

Human

Reacts with: Baboon, Cynomolgus monkey, Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG conju	Purified IgG conjugated to Pacific Blue - liquid	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Pacific Blue®	410	455

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml
Immunogen	Human monocytes
External Database Links	UniProt: P12314 Related reagents
	Entrez Gene:  2209 FCGR1A Related reagents
Synonyms	FCG1, FCGR1, IGFR1
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0-Ag14 myeloma cell line
Specificity	<b>Mouse anti Human CD64 antibody, clone 10.1</b> recognizes the human CD64 cell surface antigen, a ~75 kDa glycoprotein expressed by monocytes. The antigen acts as a high affinity receptor for human IgG, and is also known as FcRI.
	Mouse anti Human CD64 antibody, clone 10.1 blocks binding of immunoglobulin to FcRI.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul
References	1. Dougherty, G.J. <i>et al.</i> (1987) The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, 10.1. <u>Eur J Immunol. 17 (10): 1453-9.</u> 2. Beekman, J.M. <i>et al.</i> (2004) Direct interaction between FcgammaRI (CD64) and periplakin controls receptor endocytosis and ligand binding capacity. <u>Proc Natl Acad Sci U S A.101: 10392-7.</u> 3. Tanaka, M. <i>et al.</i> (2009) Activation of Fc gamma RI on monocytes triggers differentiation into immature dendritic cells that induce autoreactive T cell responses. <u>J Immunol. 183: 2349-55.</u> 4. Fet, N.G. <i>et al.</i> (2012) Reduction of activated macrophages after ischaemia-reperfusion injury diminishes oxidative stress and ameliorates renal damage. <u>Nephrol Dial Transplant.</u> 27 (8): 3149-55.  5. Wagner, C. <i>et al.</i> (2008) T lymphocytes in acute bacterial infection: increased prevalence of CD11b(+) cells in the peripheral blood and recruitment to the infected site.

- prevalence of CD11b(+) cells in the peripheral blood and recruitment to the infected site. <a href="mmunology.125:503-9">lmmunology. 125: 503-9</a>.

  6. Eisenhardt, S.U. *et al.* (2009) Dissociation of pentameric to monomeric C-reactive
- 6. Eisenhardt, S.U. *et al.* (2009) Dissociation of pentameric to monomeric C-reactive protein on activated platelets localizes inflammation to atherosclerotic plaques. <u>Circ Res.</u> 105: 128-37.
- 7. Fadlon,. E. et al. (1998) Blood polymorphonuclear leukocytes from the majority of sickle

- cell patients in the crisis phase of the disease show enhanced adhesion to vascular endothelium and increased expression of CD64. Blood. 91: 266-74.
- 8. Scheinecker, C. *et al.* (1998) Initiation of the autologous mixed lymphocyte reaction requires the expression of costimulatory molecules B7-1 and B7-2 on human peripheral blood dendritic cells. J Immunol. 161: 3966-73.
- 9. Navarro-López, F. *et al.* (2003) Late T-lymphocyte and monocyte activation in coronary restenosis. Evidence for a persistent inflammatory/immune mechanism? Rev Esp Cardiol. 56: 465-72.
- 10. Liu M *et al.* (2011) Vitellogenin mediates phagocytosis through interaction with FcγR. Mol Immunol. 49 (1-2): 211-8.
- 11. Kapelski S *et al.* (2014) Assessment of the neutrophilic antibody-dependent respiratory burst (ADRB) response to *Plasmodium falciparum*. J Leukoc Biol. 96 (6): 1131-42.
- 12. Loi, A.L.T. *et al.* (2017) Proteomic profiling of peripheral blood neutrophils identifies two inflammatory phenotypes in stable COPD patients. Respir Res. 18 (1): 100.
- 13. Hristodorov, D. *et al.* (2016) Fully human MAP-fusion protein selectively targets and eliminates proliferating CD64(+) M1 macrophages. <u>Immunol Cell Biol. 94 (5): 470-8.</u>
- 14. Kahn, F. *et al.* (2008) Antibodies against a surface protein of Streptococcus pyogenes promote a pathological inflammatory response. <u>PLoS Pathog. 4</u> (9): e1000149.

#### **Further Reading**

1. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. Exp Anim. 49 (2): 97-110.

#### Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use

#### Guarantee

12 months from date of despatch

#### Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

# Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA756PB">https://www.bio-rad-antibodies.com/SDS/MCA756PB</a> 10041

# **Related Products**

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: Pacific Blue® (MCA928PB)

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +44 (0)1865 852 739
Email: antibody\_sales\_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M344346:190111'

### Printed on 17 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody\_sales\_us@bio-rad.com