

Datasheet: MCA756SBB700

BATCH NUMBER 100008093

Description:	MOUSE ANTI HUMAN CD64:StarBright Blue 700
Specificity:	CD64
Other names:	FcRI
Format:	StarBright Blue 700
Product Type:	Monoclonal Antibody
Clone:	10.1
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Baboon, Cynomolgus monkey, Rhesus Monkey
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to StarBright Blue 700 - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
StarBright Blue 700	473	703

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Immunogen	Human monocytes
External Database Links	<p>UniProt: P12314 Related reagents</p> <p>Entrez Gene: 2209 FCGR1A Related reagents</p>
Synonyms	FCG1, FCGR1, IGFR1
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0-Ag14 myeloma cell line
Specificity	<p>Mouse anti Human CD64 antibody, clone 10.1 recognizes the human CD64 cell surface antigen, a ~75 kDa glycoprotein expressed by monocytes. The antigen acts as a high affinity receptor for human IgG, and is also known as FcRI.</p> <p>Mouse anti Human CD64 antibody, clone 10.1 blocks binding of immunoglobulin to FcRI.</p>
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	<ol style="list-style-type: none"> Scheinecker, C. <i>et al.</i> (1998) Initiation of the autologous mixed lymphocyte reaction requires the expression of costimulatory molecules B7-1 and B7-2 on human peripheral blood dendritic cells. J Immunol. 161: 3966-73. Fadlon, E. <i>et al.</i> (1998) Blood polymorphonuclear leukocytes from the majority of sickle cell patients in the crisis phase of the disease show enhanced adhesion to vascular endothelium and increased expression of CD64. Blood. 91: 266-74. Navarro-López, F. <i>et al.</i> (2003) Late T-lymphocyte and monocyte activation in coronary restenosis. Evidence for a persistent inflammatory/immune mechanism? Rev Esp Cardiol. 56: 465-72. Beekman, J.M. <i>et al.</i> (2004) Direct interaction between FcγRI (CD64) and periplakin controls receptor endocytosis and ligand binding capacity. Proc Natl Acad Sci U S A. 101: 10392-7. Kahn, F. <i>et al.</i> (2008) Antibodies against a surface protein of <i>Streptococcus pyogenes</i> promote a pathological inflammatory response. PLoS Pathog. 4 (9): e1000149. Wagner, C. <i>et al.</i> (2008) T lymphocytes in acute bacterial infection: increased prevalence of CD11b(+) cells in the peripheral blood and recruitment to the infected site.

[Immunology. 125: 503-9.](#)

7. Eisenhardt, S.U. *et al.* (2009) Dissociation of pentameric to monomeric C-reactive protein on activated platelets localizes inflammation to atherosclerotic plaques. [Circ Res. 105: 128-37.](#)
8. Tanaka, M. *et al.* (2009) Activation of Fc gamma RI on monocytes triggers differentiation into immature dendritic cells that induce autoreactive T cell responses. [J Immunol. 183: 2349-55.](#)
9. Liu M *et al.* (2011) Vitellogenin mediates phagocytosis through interaction with FcγR. [Mol Immunol. 49 \(1-2\): 211-8.](#)
10. Fet, N.G. *et al.* (2012) Reduction of activated macrophages after ischaemia-reperfusion injury diminishes oxidative stress and ameliorates renal damage. [Nephrol Dial Transplant. 27 \(8\): 3149-55.](#)
11. Kapelski S *et al.* (2014) Assessment of the neutrophilic antibody-dependent respiratory burst (ADRB) response to *Plasmodium falciparum*. [J Leukoc Biol. 96 \(6\): 1131-42.](#)
12. Hristodorov, D. *et al.* (2016) Fully human MAP-fusion protein selectively targets and eliminates proliferating CD64(+) M1 macrophages. [Immunol Cell Biol. 94 \(5\): 470-8.](#)
13. Loi, A.L.T. *et al.* (2017) Proteomic profiling of peripheral blood neutrophils identifies two inflammatory phenotypes in stable COPD patients. [Respir Res. 18 \(1\): 100.](#)
14. Rahabi, M. *et al.* (2020) Divergent Roles for Macrophage C-type Lectin Receptors, Dectin-1 and Mannose Receptors, in the Intestinal Inflammatory Response. [Cell Rep. 30 \(13\): 4386-98.e5.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA756SBB700 20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M421948:230815'

Printed on 29 Apr 2024

