

Datasheet: MCA771GT

Description:	RAT ANTI MOUSE Ly-6B.2 ALLOANTIGEN
Specificity:	Ly-6B.2 ALLOANTIGEN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	7/4
Isotype:	IgG2a
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Cultured bone marrow cells
RRID	AB_1102794
Fusion Partners	Spleen cells from AO rats were fused with cells from the Y3 Ag1.2.3 rat myeloma cell line.
Specificity	<p>Rat anti Mouse Ly-6B.2 monoclonal antibody, clone 7/4 recognizes the Ly-6B.2 antigen. Ly-6B.2 is a ~25-30 kDa GPI-anchored, heavily glycosylated protein expressed on neutrophils, inflammatory monocytes and some activated macrophages (Rosas et al. 2010). High levels of expression are seen in bone marrow, spleen, lung and lymph nodes. N-glycanase treatment of thioglycollate elicited peritoneal neutrophil lysates lowers the apparent molecular weight of Ly-6B.2 to ~15 kDa (Rosas et al.2010).</p> <p>In common with other Ly-6 antigens Ly-6B.2 demonstrates a polymorphic expression on inbred mouse strains (Kimura et al. 1984). Rat anti mouse Ly-6B.2, clone 7/4 recognizes the Ly-6B.2 antigen in 129J; AKR; C57BL/6; C57BL/10; C58; DBA/2; NZB; NZW; SJL; MFI; Swiss (PO) Strains whilst A2G; A/Sn; ASW; BALB/c; C3H/HEH; CBA.T6T6 are negative or demonstrate very weak reactivity (Hirsch and Gordon 1982).</p> <p>Rat anti mouse Ly-6B.2 has been successfully used for the immunomagnetic depletion of neutrophils during the enrichment of primitive hematopoietic cells from bone marrow (Bertoncello et al. 1991) and the depletion of myeloid cells <i>in vivo</i> (Rosas et al. 2010).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Rosas, M. <i>et al.</i> (2010) The myeloid 7/4-antigen defines recently generated inflammatory macrophages and is synonymous with Ly-6B. J Leukoc Biol. 88 (1): 169-80. Gordon, S. <i>et al.</i> (1992) Antigen markers of macrophage differentiation in murine tissues. Curr Top Microbiol Immunol. 181: 1-37. Word, R.A. <i>et al.</i> (2005) Transgene insertion on mouse chromosome 6 impairs function of the uterine cervix and causes failure of parturition. Biol Reprod. 73 (5): 1046-56. Locke, L.W. (2009) A novel neutrophil-specific PET imaging agent: cFLFLFK-PEG-64Cu. J Nucl Med. 50: 790-7. Allam, R. <i>et al.</i> (2011) Cyclic Polypeptide and Aminoglycoside Antibiotics Trigger IL-1{beta} Secretion by Activating the NLRP3 Inflammasome. J Immunol. 186: 2714-8. Holt, R. <i>et al.</i> (2011) The Molecular Mechanisms of Cervical Ripening Differ between Term and Preterm Birth. Endocrinology. 152: 1036-46. Frossard, J.L. <i>et al.</i> (2011) Role of CCL-2, CCR-2 and CCR-4 in cerulein-induced acute pancreatitis and pancreatitis-associated lung injury. J Clin Pathol. 64: 387-93. McDonald, J.U. <i>et al.</i> (2011) <i>In vivo</i> functional analysis and genetic modification of in vitro-derived mouse neutrophils. FASEB J. 25: 1972-82. Larmonier, C.B. <i>et al.</i> (2011) NHE3 modulates the severity of colitis in IL-10-deficient mice. Am J Physiol Gastrointest Liver Physiol. 300: G998-G1009. Yellon, S.M. <i>et al.</i> (2011) Remodeling of the cervix and parturition in mice lacking the progesterone receptor B isoform. Biol Reprod. 85: 498-502.

11. Nadeau, S. *et al.* (2011) Functional Recovery after Peripheral Nerve Injury is Dependent on the Pro-Inflammatory Cytokines IL-1{beta} and TNF: Implications for Neuropathic Pain. [J Neurosci. 31: 12533-12542.](#)
12. Bombardelli, L. *et al.* (2010) Pancreas-specific ablation of beta1 integrin induces tissue degeneration by disrupting acinar cell polarity. [Gastroenterology. 138: 2531-40.](#)
13. Guerriero, J.L. *et al.* (2011) DNA alkylating therapy induces tumor regression through an HMGB1-mediated activation of innate immunity. [J Immunol. 186: 3517-26.](#)
14. Brulhart-Meynet, M.C. *et al.* (2015) Improving Reconstituted HDL Composition for Efficient Post-Ischemic Reduction of Ischemia Reperfusion Injury. [PLoS One. 10 \(3\): e0119664.](#)
15. Patel, J. *et al.* (2015) RGS1 regulates myeloid cell accumulation in atherosclerosis and aortic aneurysm rupture through altered chemokine signalling. [Nat Commun. 6: 6614.](#)
16. Hamers, A.A. *et al.* (2015) Deficiency of Nuclear Receptor Nur77 Aggravates Mouse Experimental Colitis by Increased NFkB Activity in Macrophages. [PLoS One. 10 \(8\): e0133598.](#)
17. Ao, M. *et al.* (2015) Dental Infection of *Porphyromonas gingivalis* Induces Preterm Birth in Mice. [PLoS One. 10 \(8\): e0137249.](#)
18. Cnops, J. *et al.* (2015) NK-, NKT- and CD8-Derived IFN γ Drives Myeloid Cell Activation and Erythrophagocytosis, Resulting in Trypanosomiasis-Associated Acute Anemia. [PLoS Pathog. 11 \(6\): e1004964.](#)
19. Cardenas, H. *et al.* (2016) Dietary Apigenin Exerts Immune-Regulatory Activity in Vivo by Reducing NF- κ B Activity, Halting Leukocyte Infiltration and Restoring Normal Metabolic Function. [Int J Mol Sci. 17 \(3\). pii: E323. doi: 10.3390/ijms17030323.](#)
20. Park, S.W. *et al.* (2009) Human activated protein C attenuates both hepatic and renal injury caused by hepatic ischemia and reperfusion injury in mice. [Kidney Int. 76 \(7\): 739-50.](#)
21. Boal, F. *et al.* (2016) PI3P Triggers ICAM-1 Degradation in *Shigella* Infected Cells, Thus Dampening Immune Cell Recruitment. [Cell Rep. 14 \(4\): 750-9.](#)
22. Nguyen, H.T. & Shen, H. (2016) The effect of PEGylation on the stimulation of IL-1 β by gold (Au) nanoshell/silica core nanoparticles. [J Mater Chem B Mater Biol Med. 4 \(9\): 1650-1659.](#)
23. Choi, E. *et al.* (2015) Expression of Activated Ras in Gastric Chief Cells of Mice Leads to the Full Spectrum of Metaplastic Lineage Transitions. [Gastroenterology. Dec 8. pii: S0016-5085\(15\)01735-7. \[Epub ahead of print\]](#)
24. Zhao, Y. *et al.* (2016) Rapamycin prevents bronchiolitis obliterans through increasing infiltration of regulatory B cells in a murine tracheal transplantation model. [J Thorac Cardiovasc Surg. 151 \(2\): 487-496.e3.](#)
25. Wan, W. *et al.* (2015) Atypical chemokine receptor 1 deficiency reduces atherogenesis in ApoE-knockout mice. [Cardiovasc Res. 106 \(3\): 478-87.](#)
26. Kidoya, H. *et al.* (2015) APJ Regulates Parallel Alignment of Arteries and Veins in the Skin. [Dev Cell. 33 \(3\): 247-59.](#)
27. Wieser, V. *et al.* (2016) Lipocalin 2 drives neutrophilic inflammation in alcoholic liver disease. [J Hepatol. 64 \(4\): 872-80.](#)
28. Kjellman, P. *et al.* (2015) Size-dependent lymphatic uptake of nanoscale-tailored particles as tumor mass increases [Future Science OA. 1: \(4\)](#)
29. Brennan, F.H. *et al.* (2015) The Complement Receptor C5aR Controls Acute Inflammation and Astroglialosis following Spinal Cord Injury. [J Neurosci. 35 \(16\): 6517-31.](#)

30. Farrar, C.A. *et al.* (2016) Collectin-11 detects stress-induced L-fucose pattern to trigger renal epithelial injury. [J Clin Invest. Apr 18. pii: 83000. \[Epub ahead of print\]](#)
31. Laubitz, D. *et al.* (2016) Reduced Epithelial Na⁺/H⁺ Exchange Drives Gut Microbial Dysbiosis and Promotes Inflammatory Response in T Cell-Mediated Murine Colitis. [PLoS One. 11 \(4\): e0152044.](#)
32. Rabadi, M.M. *et al.* (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. [Am J Physiol Renal Physiol. Jun 22: ajprenal.00254.2016. \[Epub ahead of print\]](#)
33. Morris, A. H. *et al.* (2016) Inadequate Processing of Decellularized Dermal Matrix Reduces Cell Viability *In Vitro* and Increases Apoptosis and Acute Inflammation *In Vivo*. [BioResearch Open Access. 5 \(1\): 177-87.](#)
34. Ni, H.M. *et al.* (2016) Caspase Inhibition Prevents Tumor Necrosis Factor- α -Induced Apoptosis and Promotes Necrotic Cell Death in Mouse Hepatocytes *in Vivo* and *in Vitro*. [Am J Pathol. Sep 7. pii: S0002-9440\(16\)30236-X. \[Epub ahead of print\]](#)
35. Wang, S. *et al.* (2016) Increased hepatic receptor interacting protein kinase 3 expression due to impaired proteasomal functions contributes to alcohol-induced steatosis and liver injury. [Oncotarget. 7 \(14\): 17681-98.](#)
36. Konrad, F.M. *et al.* (2019) How Adhesion Molecule Patterns Change While Neutrophils Traffic through the Lung during Inflammation. [Mediators Inflamm. 2019: 1208086.](#)
37. Jung, P.E. *et al.* (2020) Honokiol Protects the Kidney from Renal Ischemia and Reperfusion Injury by Upregulating the Glutathione Biosynthetic Enzymes. [Biomedicines. 8 \(9\): 352.](#)
38. Chatterjee, S. *et al.* (2013) Junctional adhesion molecule-A regulates vascular endothelial growth factor receptor-2 signaling-dependent mouse corneal wound healing. [PLoS One. 8 \(5\): e63674.](#)
39. Han, Y. *et al.* (2020) Close Homolog of L1 Deficiency Exacerbated Intestinal Epithelial Barrier Function in Mouse Model of Dextran Sulfate Sodium-Induced Colitis. [Front Physiol. 11: 584508.](#)
40. Leinweber, J. *et al.* (2021) Elastase inhibitor agaphelin protects from acute ischemic stroke in mice by reducing thrombosis, blood-brain barrier damage, and inflammation. [Brain Behav Immun. S0889-1591\(20\)32485-5.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...) [FITC](#)

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos., Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®800
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M368939:200529'

Printed on 12 Feb 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)