

Datasheet: MCA772

BATCH NUMBER 165251

Description:	MOUSE ANTI RAT CD3
Specificity:	CD3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1F4
Isotype:	IgM
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/25
Immunohistology - Frozen	▪			1/10 - 1/25
Immunohistology - Paraffin (1)	▪			1/10
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **This clone is suitable for use on paraffin embedded material using target unmasking fluid [HIS003B](#), refer to [McKechnie N.M. et al.](#) for details.**

Target Species	Rat
Product Form	Purified IgM - liquid
Preparation	Purified IgM prepared by ammonium sulphate precipitation from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Approx. Protein Concentrations	IgM concentration 1.0 mg/ml
Immunogen	F344 rat T cells stimulated with PMA (TPA) and calcium ionophore
External Database Links	<p>UniProt:</p> <p>P19377 Related reagents</p> <p>Q64159 Related reagents</p> <p>Entrez Gene:</p> <p>25710 Cd3d Related reagents</p> <p>300678 Cd3g Related reagents</p>
Synonyms	T3d
RRID	AB_321258
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the P3-X63-Ag8.653 mouse myeloma cell line.
Specificity	<p>Mouse anti Rat CD3 antibody, clone 1F4 recognizes rat CD3, a ~25 kDa antigen which is found on rat T-cells. Mouse anti Rat CD3, clone 1F4 does not react with rat B cells. In immunohistology it stains rat thymus tissues strongly in the medulla and weakly in the cortex.</p> <p>Functionally the addition of the antibody to a culture of rat T cells induces the proliferation of T-cells in the presence of PMA.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Nicolls, M.G. <i>et al.</i> (1992) Induction of long-term specific tolerance to allografts in rats by therapy with an anti-CD3-like monoclonal antibody. Transplantation 55: 459-68. McKechnie NM <i>et al.</i> (1997) Immunization with the cross-reactive antigens Ov39 from <i>Onchocerca volvulus</i> and hr44 from human retinal tissue induces ocular pathology and activates retinal microglia. J Infect Dis. 176 (5): 1334-43. Candolfi, M. <i>et al.</i> (2007) Intracranial glioblastoma models in preclinical neuro-oncology: neuropathological characterization and tumor progression. J Neurooncol. 85: 133-48. Lohwasser, C. <i>et al.</i> (2009) Role of the receptor for advanced glycation end products in hepatic fibrosis. World J Gastroenterol. 15: 5789-98. Sanchez-Guajardo, V. <i>et al.</i> (2010) Microglia acquire distinct activation profiles depending on the degree of alpha-synuclein neuropathology in a rAAV based model of Parkinson's disease. PLoS One. 5: e8784. Beck, K.D. <i>et al.</i> (2010) Quantitative analysis of cellular inflammation after traumatic spinal cord injury: evidence for a multiphasic inflammatory response in the acute to chronic environment. Brain. 133: 433-47. Echeverry, S. <i>et al.</i> (2011) Peripheral Nerve Injury Alters Blood-Spinal Cord Barrier Functional and Molecular Integrity through a Selective Inflammatory Pathway. J Neurosci.

[31: 10819-28.](#)

8. Takahashi, Y. *et al.* (2017) Rituximab protects podocytes and exerts anti-proteinuric effects in rat adriamycin-induced nephropathy independent of B-lymphocytes. [Nephrology \(Carlton\). 22 \(1\): 49-57.](#)

9. Sun, J. *et al.* (2017) Pentapeptide PLNPK ameliorates adjuvant arthritis and inhibits T cell activation by suppressing Lck and PI3K activities [Int J Clin Exp Pathol 10\(5\): 5252-62.](#)

10. Du, K. *et al.* (2023) Pathogenesis of selective damage of granule cell layer in cerebellum of rats exposed to methylmercury [J Toxicolog Sci. 48 \(7\): 429-39.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA772>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) [Alk. Phos.](#)

Human Anti Mouse IgM (HCA040...) [FITC](#), [HRP](#)

Goat Anti Mouse IgM (102001...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M381503:210512'

Printed on 12 Dec 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)