

Datasheet: MCA773

BATCH NUMBER 158941

Description:	MOUSE ANTI RAT CD54
Specificity:	CD54
Other names:	ICAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1A29
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This clone is suitable for use on paraffin embedded material using target unmasking fluid [HIS003B](#) for this purpose.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Rat Ax cells (a HEV derived cell line).
External Database Links	<p>UniProt: Q00238 Related reagents</p> <p>Entrez Gene: 25464 Icam1 Related reagents</p>
Synonyms	Icam-1
RRID	AB_2280011
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells from the PAI mouse myeloma cell line.
Specificity	<p>Mouse anti Rat CD54 antibody, clone 1A29 recognizes the rat CD54 cell surface antigen, also known as intercellular adhesion molecule-1 (ICAM-1), a ~90 kDa adhesion molecule belonging to the immunoglobulin superfamily.</p> <p>CD54 is a cell surface ligand of the lymphocyte integrin, LFA-1 and plays an important role in various cell-cell interactions in the immune system. Cross-linking of ICAM-1 using clone 1A29 induces calcium signaling (Etienne <i>et al.</i> 1998).</p> <p>Mouse anti Rat CD54 antibody, clone 1A29 inhibits homotypic aggregation of cells including PHA blasts (Tamatani & Miyasaka 1990).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Etienne, S. <i>et al.</i> (1998) ICAM-1 signaling pathways associated with Rho activation in microvascular brain endothelial cells. J Immunol. 161 (10): 5755-61. Kawai, T. <i>et al.</i> (1999) Selective diapedesis of Th1 cells induced by endothelial cell RANTES. J Immunol. 163: 3269-78. Adamson, P. <i>et al.</i> (1999) Lymphocyte migration through brain endothelial cell monolayers involves signaling through endothelial ICAM-1 via a rho-dependent pathway. J Immunol. 162: 2964-73. Sato, N. <i>et al.</i> (2000) Roles of ICAM-1 for abnormal leukocyte recruitment in the microcirculation of bleomycin-induced fibrotic lung injury. Am J Respir Crit Care Med. 161: 1681-8. Etienne-Manneville, S. <i>et al.</i> (2000) ICAM-1-coupled cytoskeletal rearrangements and transendothelial lymphocyte migration involve intracellular calcium signaling in brain

- endothelial cell lines. [J Immunol. 165 \(6\): 3375-83.](#)
6. Arsenović-Ranin, N. *et al.* (2000) A monoclonal antibody to the rat Crry/p65 antigen, a complement regulatory membrane protein, stimulates adhesion and proliferation of thymocytes. [Immunology. 100: 334-44.](#)
 7. Beck-Schimmer, B. *et al.* (2001) Hypoxia mediates increased neutrophil and macrophage adhesiveness to alveolar epithelial cells. [Am J Respir Cell Mol Biol. 25: 780-7.](#)
 8. McKechnie, N. M. *et al.* (2002) Antigenic mimicry: *Onchocerca volvulus* antigen-specific T cells and ocular inflammation. [Invest Ophthalmol Vis Sci. 43:411-8.](#)
 9. Zhu, X. *et al.* (2003) Matrine protects sinusoidal endothelial cells from cold ischemia and reperfusion injury in rat orthotopic liver transplantation. [Ann Clin Lab Sci. 33: 216-25.](#)
 10. Deng, H. *et al.* (2003) Mild hypothermia inhibits inflammation after experimental stroke and brain inflammation. [Stroke. 34: 2495-501.](#)
 11. Ikezumi, Y. *et al.* (2004) Macrophage-mediated renal injury is dependent on signaling via the JNK pathway. [J Am Soc Nephrol. 15: 1775-84.](#)
 12. Westermann, J. *et al.* (2005) Naive, effector, and memory T lymphocytes efficiently scan dendritic cells *in vivo*: contact frequency in T cell zones of secondary lymphoid organs does not depend on LFA-1 expression and facilitates survival of effector T cells. [J Immunol. 174: 2517-24.](#)
 13. Westermann, D. *et al.* (2007) Cardioprotective and anti-inflammatory effects of interleukin converting enzyme inhibition in experimental diabetic cardiomyopathy. [Diabetes. 56: 1834-41.](#)
 14. Couty, J.P. (2007) PECAM-1 engagement counteracts ICAM-1-induced signaling in brain vascular endothelial cells. [J Neurochem. 103: 793-801.](#)
 15. Trinh, L. *et al.* (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between *in vivo* confocal microscopy and immunohistochemistry. [Mol Vis. 14: 1149-56.](#)
 16. Kanellis, J. *et al.* (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. [Nephrol Dial Transplant. 25: 2898-908.](#)
 17. Azcutia V *et al.* (2010) Inflammation determines the pro-adhesive properties of high extracellular d-glucose in human endothelial cells *in vitro* and rat microvessels *in vivo*. [PLoS One. 5 \(4\): e10091.](#)
 18. Choi, J.S. *et al.* (2011) Mild Hypothermia Attenuates Intercellular Adhesion Molecule-1 Induction via Activation of Extracellular Signal-Regulated Kinase-1/2 in a Focal Cerebral Ischemia Model. [Stroke Res Treat. 2011: 846716.](#)
 19. Li, W. & Klein, S.L. (2012) Seoul virus-infected rat lung endothelial cells and alveolar macrophages differ in their ability to support virus replication and induce regulatory T cell phenotypes. [J Virol. 86 \(21\): 11845-55.](#)
 20. Gates, D. *et al.* (2012) Apo J/clusterin expression and secretion: evidence for 15-deoxy- $\Delta(12,14)$ -PGJ(2)-dependent mechanism. [Biochim Biophys Acta. 1821 \(2\): 335-42.](#)
 21. Liu, Y.C. *et al.* (2013) A biodegradable, sustained-released, prednisolone acetate microfilm drug delivery system effectively prolongs corneal allograft survival in the rat keratoplasty model. [PLoS One. 8 \(8\): e70419.](#)
 22. Li, Z. *et al.* (2015) Three-dimensional graphene foams loaded with bone marrow derived mesenchymal stem cells promote skin wound healing with reduced scarring. [Mater Sci Eng C Mater Biol Appl. 57: 181-8.](#)

23. Gautier, S. *et al.* (2015) PPAR-Alpha Agonist Used at the Acute Phase of Experimental Ischemic Stroke Reduces Occurrence of Thrombolysis-Induced Hemorrhage in Rats. [PPAR Res. 2015: 246329.](#)
24. Ichihara, Y. *et al.* (2018) Self-assembling peptide hydrogel enables instant epicardial coating of the heart with mesenchymal stromal cells for the treatment of heart failure. [Biomaterials. 154: 12-23.](#)
25. Cakała-Jakimowicz, M. & Puzianowska-Kuznicka, M. (2022) Towards Understanding the Lymph Node Response to Skin Infection with Saprophytic *Staphylococcus epidermidis*. [Biomedicines. 10 \(5\): 1021.](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA773 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Human Anti Mouse IgG1 (HCA036...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M387350:210623'

