

Datasheet: MCA773

| | |
|----------------------|---------------------|
| Description: | MOUSE ANTI RAT CD54 |
| Specificity: | CD54 |
| Other names: | ICAM-1 |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | 1A29 |
| Isotype: | IgG1 |
| Quantity: | 0.2 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/25 - 1/50 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin (1) | ▪ | | | |
| ELISA | | | ▪ | |
| Immunoprecipitation | ▪ | | | |
| Western Blotting | ▪ | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **This clone is suitable for use on paraffin embedded material using target unmasking fluid [HIS003B](#) for this purpose.**

| | |
|------------------------|---|
| Target Species | Rat |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |

| | |
|---------------------------------------|---|
| Preservative Stabilisers | 0.09% Sodium Azide |
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |
| Immunogen | Rat Ax cells (a HEV derived cell line). |
| External Database Links | <p>UniProt: Q00238 Related reagents</p> <p>Entrez Gene: 25464 Icam1 Related reagents</p> |
| Synonyms | Icam-1 |
| RRID | AB_2280011 |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells from the PAI mouse myeloma cell line. |
| Specificity | <p>Mouse anti Rat CD54 antibody, clone 1A29 recognizes the rat CD54 cell surface antigen, also known as intercellular adhesion molecule-1 (ICAM-1), a ~90 kDa adhesion molecule belonging to the immunoglobulin superfamily.</p> <p>CD54 is a cell surface ligand of the lymphocyte integrin, LFA-1 and is known to play an important role in various cell-cell interactions in the immune system. Studies suggest that cross-linking of ICAM-1 using clone 1A29 induces calcium signalling (Etienne <i>et al.</i>).</p> <p>Functionally Mouse anti Rat CD54 antibody, clone 1A29 inhibits homotypic aggregation of PHA blasts.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| References | <ol style="list-style-type: none"> 1. Kawai, T. <i>et al.</i> (1999) Selective diapedesis of Th1 cells induced by endothelial cell RANTES. J Immunol. 163: 3269-78. 2. Sato, N. <i>et al.</i> (2000) Roles of ICAM-1 for abnormal leukocyte recruitment in the microcirculation of bleomycin-induced fibrotic lung injury. Am J Respir Crit Care Med. 161: 1681-8. 3. Etienne, S. <i>et al.</i> (1998) ICAM-1 signaling pathways associated with Rho activation in microvascular brain endothelial cells. J Immunol. 161 (10): 5755-61. 4. Etienne-Manneville, S. <i>et al.</i> (2000) ICAM-1-coupled cytoskeletal rearrangements and transendothelial lymphocyte migration involve intracellular calcium signaling in brain endothelial cell lines. J Immunol. 165 (6): 3375-83. 5. McKechnie, N. M. <i>et al.</i> (2002) Antigenic mimicry: Onchocerca volvulus antigen-specific T cells and ocular inflammation. Invest Ophthalmol Vis Sci. 43:411-8. |

6. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. [Br J Ophthalmol. 2003 Dec;87\(12\):1515-22.](#)
7. Kielian, T. *et al.* (2000) Proinflammatory cytokine, chemokine, and cellular adhesion molecule expression during the acute phase of experimental brain abscess development. [Am J Pathol. 157: 647-58.](#)
8. Trinh, L. *et al.* (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between *in vivo* confocal microscopy and immunohistochemistry. [Mol Vis. 14: 1149-56.](#)
9. Adamson, P. *et al.* (1999) Lymphocyte migration through brain endothelial cell monolayers involves signaling through endothelial ICAM-1 via a rho-dependent pathway. [J Immunol. 162: 2964-73.](#)
10. Arsenović-Ranin, N. *et al.* (2000) A monoclonal antibody to the rat Crry/p65 antigen, a complement regulatory membrane protein, stimulates adhesion and proliferation of thymocytes. [Immunology. 100: 334-44.](#)
11. Beck-Schimmer, B. *et al.* (2001) Hypoxia mediates increased neutrophil and macrophage adhesiveness to alveolar epithelial cells. [Am J Respir Cell Mol Biol. 25: 780-7.](#)
12. Couty, J.P. (2007) PECAM-1 engagement counteracts ICAM-1-induced signaling in brain vascular endothelial cells. [J Neurochem. 103: 793-801.](#)
13. Deng, H. *et al.* (2003) Mild hypothermia inhibits inflammation after experimental stroke and brain inflammation. [Stroke. 34: 2495-501.](#)
14. Ikezumi, Y. *et al.* (2004) Macrophage-mediated renal injury is dependent on signaling via the JNK pathway. [J Am Soc Nephrol. 15: 1775-84.](#)
15. Kanellis, J. *et al.* (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. [Nephrol Dial Transplant. 25: 2898-908.](#)
16. Westermann, D. *et al.* (2007) Cardioprotective and anti-inflammatory effects of interleukin converting enzyme inhibition in experimental diabetic cardiomyopathy. [Diabetes. 56: 1834-41.](#)
17. Zhu, X. *et al.* (2003) Matrine protects sinusoidal endothelial cells from cold ischemia and reperfusion injury in rat orthotopic liver transplantation. [Ann Clin Lab Sci. 33: 216-25.](#)
18. Choi, J.S. *et al.* (2011) Mild Hypothermia Attenuates Intercellular Adhesion Molecule-1 Induction via Activation of Extracellular Signal-Regulated Kinase-1/2 in a Focal Cerebral Ischemia Model. [Stroke Res Treat. 2011: 846716.](#)
19. Azcutia V *et al.* (2010) Inflammation determines the pro-adhesive properties of high extracellular d-glucose in human endothelial cells *in vitro* and rat microvessels *in vivo*. [PLoS One. 5 \(4\): e10091.](#)
20. Li, W. & Klein, S.L. (2012) Seoul virus-infected rat lung endothelial cells and alveolar macrophages differ in their ability to support virus replication and induce regulatory T cell phenotypes. [J Virol. 86 \(21\): 11845-55.](#)
21. Gates, D. *et al.* (2012) Apo J/clusterin expression and secretion: evidence for 15-deoxy- $\Delta(12,14)$ -PGJ(2)-dependent mechanism. [Biochim Biophys Acta. 1821 \(2\): 335-42.](#)
22. Li, Z. *et al.* (2015) Three-dimensional graphene foams loaded with bone marrow derived mesenchymal stem cells promote skin wound healing with reduced scarring. [Mater Sci Eng C Mater Biol Appl. 57: 181-8.](#)
23. Gautier, S. *et al.* (2015) PPAR-Alpha Agonist Used at the Acute Phase of Experimental Ischemic Stroke Reduces Occurrence of Thrombolysis-Induced Hemorrhage

in Rats. [PPAR Research. 2015: 1-6.](#)

24. Liu, Y.C. *et al.* (2013) A biodegradable, sustained-released, prednisolone acetate microfilm drug delivery system effectively prolongs corneal allograft survival in the rat keratoplasty model. [PLoS One. 8 \(8\): e70419.](#)

25. Ichihara, Y. *et al.* (2018) Self-assembling peptide hydrogel enables instant epicardial coating of the heart with mesenchymal stromal cells for the treatment of heart failure. [Biomaterials. 154: 12-23.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Human Anti Mouse IgG1 (HCA036...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M387350:210623'

Printed on 21 Sep 2021