

## Datasheet: MCA773GA

**BATCH NUMBER 173169**

<b>Description:</b>	MOUSE ANTI RAT CD54
<b>Specificity:</b>	CD54
<b>Other names:</b>	ICAM-1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	1A29
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **This clone is suitable for use on paraffin embedded material following antigen retrieval ([McKechnie N.M. et al. 1997](#)).**

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Rat Ax cells (a HEV derived cell line).
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q00238</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">25464</a>    Icam1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Icam-1
<b>RRID</b>	AB_566847
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells from the PAI mouse myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat CD54 antibody, clone 1A29</b> recognizes the rat CD54 cell surface antigen, also known as intercellular adhesion molecule-1 (ICAM-1), a ~90 kDa adhesion molecule belonging to the immunoglobulin superfamily.</p> <p>CD54 is a cell surface ligand of the lymphocyte integrin, LFA-1 and plays an important role in various cell-cell interactions in the immune system. Cross-linking of ICAM-1 using clone 1A29 induces calcium signaling (<a href="#">Etienne <i>et al.</i> 1998</a>).</p> <p>Mouse anti Rat CD54 antibody, clone 1A29 inhibits homotypic aggregation of cells including PHA blasts (<a href="#">Tamatani &amp; Miyasaka 1990</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Etienne, S. <i>et al.</i> (1998) ICAM-1 signaling pathways associated with Rho activation in microvascular brain endothelial cells. <a href="#">J Immunol. 161 (10): 5755-61.</a></li> <li>Kawai, T. <i>et al.</i> (1999) Selective diapedesis of Th1 cells induced by endothelial cell RANTES. <a href="#">J Immunol. 163: 3269-78.</a></li> <li>Adamson, P. <i>et al.</i> (1999) Lymphocyte migration through brain endothelial cell monolayers involves signaling through endothelial ICAM-1 via a rho-dependent pathway. <a href="#">J Immunol. 162: 2964-73.</a></li> <li>Sato, N. <i>et al.</i> (2000) Roles of ICAM-1 for abnormal leukocyte recruitment in the microcirculation of bleomycin-induced fibrotic lung injury. <a href="#">Am J Respir Crit Care Med. 161: 1681-8.</a></li> <li>Etienne-Manneville, S. <i>et al.</i> (2000) ICAM-1-coupled cytoskeletal rearrangements and transendothelial lymphocyte migration involve intracellular calcium signaling in brain</li> </ol>

- endothelial cell lines. [J Immunol. 165 \(6\): 3375-83.](#)
6. Arsenović-Ranin, N. *et al.* (2000) A monoclonal antibody to the rat Crry/p65 antigen, a complement regulatory membrane protein, stimulates adhesion and proliferation of thymocytes. [Immunology. 100: 334-44.](#)
7. Beck-Schimmer, B. *et al.* (2001) Hypoxia mediates increased neutrophil and macrophage adhesiveness to alveolar epithelial cells. [Am J Respir Cell Mol Biol. 25: 780-7.](#)
8. McKechnie, N. M. *et al.* (2002) Antigenic mimicry: *Onchocerca volvulus* antigen-specific T cells and ocular inflammation. [Invest Ophthalmol Vis Sci. 43:411-8.](#)
9. Zhu, X. *et al.* (2003) Matrine protects sinusoidal endothelial cells from cold ischemia and reperfusion injury in rat orthotopic liver transplantation. [Ann Clin Lab Sci. 33: 216-25.](#)
10. Deng, H. *et al.* (2003) Mild hypothermia inhibits inflammation after experimental stroke and brain inflammation. [Stroke. 34: 2495-501.](#)
11. Ikezumi, Y. *et al.* (2004) Macrophage-mediated renal injury is dependent on signaling via the JNK pathway. [J Am Soc Nephrol. 15: 1775-84.](#)
12. Westermann, J. *et al.* (2005) Naive, effector, and memory T lymphocytes efficiently scan dendritic cells *in vivo*: contact frequency in T cell zones of secondary lymphoid organs does not depend on LFA-1 expression and facilitates survival of effector T cells. [J Immunol. 174: 2517-24.](#)
13. Westermann, D. *et al.* (2007) Cardioprotective and anti-inflammatory effects of interleukin converting enzyme inhibition in experimental diabetic cardiomyopathy. [Diabetes. 56: 1834-41.](#)
14. Couty, J.P. (2007) PECAM-1 engagement counteracts ICAM-1-induced signaling in brain vascular endothelial cells. [J Neurochem. 103: 793-801.](#)
15. Trinh, L. *et al.* (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between *in vivo* confocal microscopy and immunohistochemistry. [Mol Vis. 14: 1149-56.](#)
16. Kanellis, J. *et al.* (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. [Nephrol Dial Transplant. 25: 2898-908.](#)
17. Azcutia V *et al.* (2010) Inflammation determines the pro-adhesive properties of high extracellular d-glucose in human endothelial cells *in vitro* and rat microvessels *in vivo*. [PLoS One. 5 \(4\): e10091.](#)
18. Choi, J.S. *et al.* (2011) Mild Hypothermia Attenuates Intercellular Adhesion Molecule-1 Induction via Activation of Extracellular Signal-Regulated Kinase-1/2 in a Focal Cerebral Ischemia Model. [Stroke Res Treat. 2011: 846716.](#)
19. Li, W. & Klein, S.L. (2012) Seoul virus-infected rat lung endothelial cells and alveolar macrophages differ in their ability to support virus replication and induce regulatory T cell phenotypes. [J Virol. 86 \(21\): 11845-55.](#)
20. Gates, D. *et al.* (2012) Apo J/clusterin expression and secretion: evidence for 15-deoxy- $\Delta$ (12,14)-PGJ(2)-dependent mechanism. [Biochim Biophys Acta. 1821 \(2\): 335-42.](#)
21. Liu, Y.C. *et al.* (2013) A biodegradable, sustained-released, prednisolone acetate microfilm drug delivery system effectively prolongs corneal allograft survival in the rat keratoplasty model. [PLoS One. 8 \(8\): e70419.](#)
22. Li, Z. *et al.* (2015) Three-dimensional graphene foams loaded with bone marrow derived mesenchymal stem cells promote skin wound healing with reduced scarring. [Mater Sci Eng C Mater Biol Appl. 57: 181-8.](#)

23. Gautier, S. *et al.* (2015) PPAR-Alpha Agonist Used at the Acute Phase of Experimental Ischemic Stroke Reduces Occurrence of Thrombolysis-Induced Hemorrhage in Rats. [PPAR Res. 2015: 246329.](#)
24. Ichihara, Y. *et al.* (2018) Self-assembling peptide hydrogel enables instant epicardial coating of the heart with mesenchymal stromal cells for the treatment of heart failure. [Biomaterials. 154: 12-23.](#)
25. Çakala-Jakimowicz, M. & Puzianowska-Kuznicka, M. (2022) Towards Understanding the Lymph Node Response to Skin Infection with Saprophytic *Staphylococcus epidermidis*. [Biomedicines. 10 \(5\): 1021.](#)

<b>Storage</b>	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA773GA">https://www.bio-rad-antibodies.com/SDS/MCA773GA</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Human Anti Mouse IgG1 (HCA036...)	<a href="#">HRP</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)  
'M433790:241212'

Printed on 19 Mar 2026

