

# Datasheet: MCA796GA

**BATCH NUMBER 152472**

<b>Description:</b>	MOUSE ANTI HUMAN CD62P
<b>Specificity:</b>	CD62P
<b>Other names:</b>	P-SELECTIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AK-6
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen	▪			1/1000 - 1/5000
Immunohistology - Paraffin (1)	▪			1/500 - 1/2000
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1) Staining may be enhanced using Sodium Citrate buffer pH6.0**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Rhesus Monkey</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human platelet membrane glycoproteins.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P16109</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">6403</a>    SELP    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	GMRP, GRMP
<b>RRID</b>	AB_1125272
<b>Specificity</b>	<p><b>Mouse anti Human CD62P antibody, clone AK-6</b> recognizes the CD62P, also known as P-selectin, Granule membrane protein 140, GMP140, Leukocyte-endothelial cell adhesion molecule 3 or Platelet activation dependent granule-external membrane protein. CD62P is a 830 amino acid, including a 41 amino acid signal peptide, ~140 kDa single pass type I transmembrane glycoprotein expressed on activated platelets and endothelial cell</p> <p>CD62P plays an important role in adhesive processes between leucocytes and endothelial cells. CD62P is a component of the platelet alpha granule and is rapidly translocated to the plasma membrane upon activation (<a href="#">Stenberg et al. 1985</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $10^6$ cells in 100ul.
<b>Immunohistology</b>	This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections but staining may be enhanced using Sodium Citrate buffer pH6.0
<b>Histology Positive Control Tissue</b>	Tonsil
<b>References</b>	<ol style="list-style-type: none"> <li>1. Dunlop, L.C. <i>et al.</i> (1992) Characterization of GMP-140 (P-selectin) as a circulating plasma protein. <a href="#">J Exp Med. 175 (4): 1147-50.</a></li> <li>2. Skinner, M.P. <i>et al.</i> (1989) Characterization of human platelet GMP-140 as a heparin-</li> </ol>

- binding protein. [Biochem Biophys Res Commun. 164 \(3\): 1373-9.](#)
3. Skinner, M.P. *et al.* (1991) GMP-140 binding to neutrophils is inhibited by sulfated glycans. [J Biol Chem. 266 \(9\): 5371-4.](#)
  4. Sopper, S. *et al.* (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. [Cytometry. 29 \(4\): 351-62.](#)
  5. Bevilacqua, M.P. & Nelson, R.M. (1993) Selectins. [J Clin Invest. 91 \(2\): 379-87.](#)
  6. Roos-Engstrand, E. *et al.* (2005) Increased expression of p38 MAPK in human bronchial epithelium after lipopolysaccharide exposure. [Eur Respir J. 25 \(5\): 797-803.](#)
  7. Kornerup, K.N. *et al.* (2010) Circulating platelet-neutrophil complexes are important for subsequent neutrophil activation and migration. [J Appl Physiol. 109: 758-67.](#)
  8. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on in vitro and in vivo lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
  9. Dalli, J. *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
  10. Wassmer, S.C. *et al.* (2008) Platelet-induced clumping of Plasmodium falciparum-infected erythrocytes from Malawian patients with cerebral malaria-possible modulation in vivo by thrombocytopenia. [J Infect Dis. 197: 72-8.](#)
  11. Theoret, J.F. *et al.* (2001) P-selectin antagonism with recombinant p-selectin glycoprotein ligand-1 (rPSGL-Ig) inhibits circulating activated platelet binding to neutrophils induced by damaged arterial surfaces. [J Pharmacol Exp Ther. 298: 658-64](#)
  12. Turner, C.P. *et al.* (2003) The role of P-selectin in the immune destruction of platelets. [Br J Haematol. 121: 623-31.](#)
  13. van Nispen tot Pannerden, H. *et al.* (2010) The platelet interior revisited: electron tomography reveals tubular alpha-granule subtypes. [Blood. 116: 1147-56.](#)
  14. Knipe, L. *et al.* (2010) A revised model for the secretion of tPA and cytokines from cultured endothelial cells. [Blood. 116 \(12\): 2183-91.](#)
  15. Kitaya, K. & Yasuo, T. (2010) Aberrant expression of selectin E, CXCL1, and CXCL13 in chronic endometritis. [Mod Pathol. 23 \(8\): 1136-46.](#)
  16. Xiong, G.M. *et al.* (2015) Imparting electroactivity to polycaprolactone fibers with heparin-doped polypyrrole: Modulation of hemocompatibility and inflammatory responses. [Acta Biomater. 23: 240-9.](#)
  17. Liao, Y. *et al.* (2017) Tailoring of TiO<sub>2</sub> films by H<sub>2</sub>SO<sub>4</sub> treatment and UV irradiation to improve anticoagulant ability and endothelial cell compatibility. [Colloids Surf B Biointerfaces. 155: 314-22.](#)
  18. Christersson, C. *et al.* (2013) Evaluation of microparticles in whole blood by multicolour flow cytometry assay. [Scand J Clin Lab Invest. 73\(3\): 229-39.](#)
  19. Tardy-Poncet, B. *et al.* (2021) Functional Flow Cytometric Assay for Reliable and Convenient Heparin-Induced Thrombocytopenia Diagnosis in Daily Practice [Biomedicines. 9 \(4\): 332.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA796GA">https://www.bio-rad-antibodies.com/SDS/MCA796GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M368977:200529'

Printed on 19 Jan 2024