

Datasheet: MCA799GA

Description:	MOUSE ANTI RABBIT CD4
Specificity:	CD4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	KEN-4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Rabbit

Species Cross Reactivity

Reacts with: Brown Hare (*Lepus europeus*)

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Rabbit thymocytes.
External Database Links	<p>UniProt: P46630 Related reagents</p> <p>Entrez Gene: 100009152 CD4 Related reagents</p>
Fusion Partners	Spleen cells from immunized mice were fused with cells of the mouse PAI myeloma cell line.
Specificity	<p>Mouse anti Rabbit CD4 antibody, clone KEN-4 recognizes the rabbit CD4 cell surface antigen, also known as T-cell surface antigen T4/Leu-3. Rabbit CD4 is a 434 amino acid, with an additional N-terminal signal peptide ~50 kDa cell surface single pass, type I transmembrane glycoprotein expressed by T helper cells.</p> <p>Mouse anti Rabbit CD4 antibody, clone KEN-4 blocks the allogeneic mixed lymphocyte reaction response.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
References	<ol style="list-style-type: none"> Perosa, F. and Dammacco, F. (1994) Anti-idiotypic monoclonal antibodies (mAb) to an anti-CD4 mAb induce CD4+ T cell depletion in rabbit. Int J Clin Lab Res. 24: 208-12. Renaux, S. <i>et al.</i> (2003) Dynamics and responsiveness of T-lymphocytes in secondary lymphoid organs of rabbits developing immunity to <i>Eimeria intestinalis</i>. Vet Parasitol. 110 (3-4): 181-95. Dewals, B. <i>et al.</i> (2008) Malignant catarrhal fever induced by alcelaphine herpesvirus 1 is associated with proliferation of CD8+ T cells supporting a latent infection. PLoS ONE 3: e1627. Pakandl, M. <i>et al.</i> (2008) Dependence of the immune response to coccidiosis on the age of rabbit suckling. Parasitol Res. 103 (6): 1265-71. Yang, J. <i>et al.</i> (2009) Expression and localization of rabbit B-cell activating factor (BAFF) and its specific receptor BR3 in cells and tissues of the rabbit immune system. Dev Comp Immunol. 33 (5): 697-708. Chentoufi, A.A. <i>et al.</i> (2010) A novel HLA (HLA-A*0201) transgenic rabbit model for preclinical evaluation of human CD8+ T cell epitope-based vaccines against ocular herpes. J Immunol. 184: 2561-71. Rütgen, B.C. <i>et al.</i> (2014) Exploratory assessment of CD4+ T lymphocytes in brown hares (<i>Lepus europeus</i>) using a cross-reactive anti-rabbit CD4 antibody. Vet Immunol

[Immunopathol. 161 \(1-2\): 108-15.](#)

8. Parameswaran, N. *et al.* (2014) The A2 gene of alcelaphine herpesvirus-1 is a transcriptional regulator affecting cytotoxicity in virus-infected T cells but is not required for malignant catarrhal fever induction in rabbits. [Virus Res. 188: 68-80.](#)
9. Boutard, B. *et al.* (2015) The α 2,3-sialyltransferase encoded by myxoma virus is a virulence factor that contributes to immunosuppression. [PLoS One. 10 \(2\): e0118806.](#)
10. Khan, A.A. *et al.* (2015) Therapeutic immunization with a mixture of herpes simplex virus 1 glycoprotein D-derived "asymptomatic" human CD8+ T-cell epitopes decreases spontaneous ocular shedding in latently infected HLA transgenic rabbits: association with low frequency of local PD-1+ TIM-3+ CD8+ exhausted T cells. [J Virol. 89 \(13\): 6619-32.](#)
11. Myster, F. *et al.* (2015) Viral semaphorin inhibits dendritic cell phagocytosis and migration but is not essential for gammaherpesvirus-induced lymphoproliferation in malignant catarrhal fever. [J Virol. 89 \(7\): 3630-47.](#)
12. Beghelli, D *et al.* (2016) Effects of Oregano (*Origanum vulgare* L.) and Rosemary (*Rosmarinus officinalis* L.) Aqueous Extracts On *in vitro* Rabbit Immune Responses [MOJ Immunology. 4 \(4\) \[Epub ahead of print\].](#)
13. Sorel, O. *et al.* (2017) Macavirus latency-associated protein evades immune detection through regulation of protein synthesis in cis depending upon its glycin/glutamate-rich domain. [PLoS Pathog. 13 \(10\): e1006691.](#)
14. Penadés, M. *et al.* (2018) Long-term implications of feed energy source in different genetic types of reproductive rabbit females. II. Immunologic status. [Animal. 12 \(9\): 1877-85.](#)
15. Jeklova, E. *et al.* (2020) Characterization of humoral and cell-mediated immunity in rabbits orally infected with *Encephalitozoon cuniculi*.. [Vet Res. 51 \(1\): 79.](#)
16. Niedźwiedzka-Rystwej, P. *et al.* (2020) B and T lymphocytes in rabbits change according to the sex and throughout the year. [Pol J Vet Sci. 23 \(1\): 37-42.](#)
17. Muñoz-Silvestre, A. *et al.* (2020) Pathogenesis of Intradermal Staphylococcal Infections: Rabbit Experimental Approach to Natural *Staphylococcus aureus* Skin Infections. [Am J Pathol. 190 \(6\): 1188-1210.](#)
18. Largo, R.D. *et al.* (2020) VEGF Over-Expression by Engineered BMSC Accelerates Functional Perfusion, Improving Tissue Density and In-Growth in Clinical-Size Osteogenic Grafts. [Front Bioeng Biotechnol. 8: 755.](#)
19. Niedźwiedzka-Rystwej, P. *et al.* (2022) Reactivity of selected markers of innate and adaptive immunity in rabbits experimentally infected with antigenic variants of RHD (Lagovirus europaeus/GI.1a). [Vet Res Commun. 46 \(1\): 233-42.](#)
20. Dewals, B.G. & Vanderplasschen, A. (2011) Malignant catarrhal fever induced by Alcelaphine herpesvirus 1 is characterized by an expansion of activated CD3+CD8+CD4- T cells expressing a cytotoxic phenotype in both lymphoid and non-lymphoid tissues. [Vet Res. 42 \(1\): 95.](#)
21. Li, H. *et al.* (2011) Characterization of ovine herpesvirus 2-induced malignant catarrhal fever in rabbits. [Vet Microbiol. 150 \(3-4\): 270-7.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in

frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA799GA 10040
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M382563:210513'

Printed on 29 Feb 2024