

Datasheet: MCA808GA BATCH NUMBER 164282

Description:	MOUSE ANTI RABBIT CD45
Specificity:	CD45
Other names:	LCA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	L12/201
lsotype:	lgG1
Quantity:	0.1 mg

Product Details

Buffer Solution

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				1/50 - 1/100
	Immunohistology - Frozen				
	Immunohistology - Paraffin			-	
	ELISA			-	
	Immunoprecipitation			-	
	Western Blotting			•	
	Immunofluorescence				
	necessarily exclude its use in such procedures. Suggested working dilution a guide only. It is recommended that the user titrates the antibody for use system using appropriate negative/positive controls.				king dilutions are given as
	a guide only. It is recomm system using appropriate	nended th negative	hat the use	er titrates the antibo controls.	dy for use in their own
Target Species	a guide only. It is recomm system using appropriate Rabbit	nended th negative	hat the use	controls.	dy for use in their own
Target Species Product Form	a guide only. It is recomm system using appropriate Rabbit Purified IgG - liquid	nended th negative	, hat the us ∳/positive	controls.	dy for use in their own

Preservative 0.09% Sodium Azide (NaN₃)

Phosphate buffered saline

Stabilisers

Carrier Free	Yes	
Approx. Protein Concentrations	IgG concentration 1.0mg/ml	
Immunogen	Glycoproteins isolated from the T cell line, RL-5.	
RRID	AB_10961760	
Fusion Partners	Spleen cells from immunised mice were fused with cells of the I myeloma cell line.	P3.X63.Ag8-U1 mouse
Specificity	Mouse anti Rabbit CD45 antibody, clone L12/201 recognizes known as leukocyte common antigen (LCA) or T200. Mouse an clone L12/201 shows pan leucocyte reactivity by flow cytometry immunohistochemistry.	s the CD45 antigen, also nti Rabbit CD45 antibody, / and
	Immunoprecipitation was achieved by cross linking antibody to yielding a protein migrating by gel electrophoresis at a molecula	the labelled cell surface ar mass of ~200 kDa.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or	cells or 100ul whole blood.
References	 Jackson, S. <i>et al.</i> (1983) Differentiation antigens identify subples lymphocytes. Definition by flow cytometry. J Exp Med. 157 (1) Wilkinson, J.M. <i>et al.</i> (1984) Cell surface glycoproteins of rabic characterization with monoclonal antibodies. Mol Immunol. 21 (Wilkinson, J.M. <i>et al.</i> (1992) A cytotoxic rabbit T-cell line infervirus which expresses CD8 and class II antigens. Immunology. Wilkinson, J.M. <i>et al.</i> (1993) Immunohistochemical identificat populations in normal tissue and inflamed synovium of the rabb 315-20. Fenton, M. <i>et al.</i> (2001) Cellular senescence after single and denudations of rabbit carotid arteries. Arterioscler Thromb Vasc 6. Mackenzie, S.M. <i>et al.</i> (2006) Immunocontraceptive effects of with recombinant myxoma virus expressing rabbit ZP2 or ZP3. 7. Davis, W.C. & Hamilton, M.J. (2008) Use of flow cytometry to a set of monoclonal antibodies specific for rabbit leukocyte differences Sci. 9 (1): 51-66. Liang, H. <i>et al.</i> (2009) Comparison of the ocular tolerability of emulsion versus conventional formulations of prostaglandins: an Vis. 15: 1690-9. Xu, Y. <i>et al.</i> (2010) Adenovirus-mediated overexpression of g mitigates transplant arteriosclerosis in rabbit carotid allografts. 1409-16. Vasíček, J <i>et al.</i> (2014) Basic Blood Analysis of Rabbits Imm Against Myxomatosis. Proc Int Symp Anim Sci 2014: 411-6. 	populations of rabbit T and): 34-46. obit lymphocytes: 1): 95-103. cted with a gamma-herpes 77 (1): 106-8. tion of leucocyte oit. J Pathol. 170 (3): I repeated balloon catheter 2 Biol. 21: 220-6. on female rabbits infected Biol Reprod. 74: 511-21. o develop and characterize erentiation molecules. J Vet f a latanoprost cationic n <i>in vivo</i> toxicity assay. Mol glutathione-s-transferase Transplantation. 89: munized with Vaccine

	11. Vasicek, J. <i>et al.</i> (2015) Determination of Lymphocyte Subset Distribution in the Peripheral Blood of Rabbits Immunized with CFA. <u>Int Symp Anim Sci UDC:639.112 pp.</u> 226-31.
	 12. Sijnave, D. <i>et al.</i> (2015) Inhibition of Rho-Associated Kinase Prevents Pathological Wound Healing and Neovascularization After Corneal Trauma. <u>Cornea. 34 (9): 1120-9.</u> 13. Ondruska, L. <i>et al.</i> (2016) Decrease in C-reactive protein levels in rabbits after vaccination with a live attenuated myxoma virus vaccine <u>Veterinární Medicína. 61 (No. 10): 571-6.</u>
	14. Ondruska, L <i>et al.</i> (2016) Decrease in C-reactive protein levels in rabbits after vaccination with a live attenuated myxoma virus vaccine <u>Veterinární Medicína. 61 (No.</u> 10): 571-6.
	 15. Kováč, M. <i>et al.</i> (2017) Phenotype and ultrastructure of stem cells derived from amniotic fluid of Nitra rabbit <u>J Cent Eur Agric.18 (1): 226-34.</u> 16. Kovac, M. <i>et al.</i> (2017) Different RNA and protein expression of surface markers in
	 rabbit amniotic fluid-derived mesenchymal stem cells. <u>Biotechnol Prog. 33 (6): 1601-13.</u> 17. Vašíček, J <i>et al.</i> (2018) The Efficiency of Immunomagnetic Sorting of Rabbit Bone Marrow Celld for the Establishment of Mesenchymal Stem Cell Culture. <u>J Microbiol</u> <u>Biotech Food Sci. 8 (3): 890-2.</u>
	 Barth, H. <i>et al.</i> (2019) Inflammatory responses after vitrectomy with vitreous substitutes in a rabbit model. <u>Graefes Arch Clin Exp Ophthalmol. 257 (4): 769-83.</u> Kulikova, B. <i>et al.</i> (2019) Survivability of rabbit amniotic fluid-derived mesenchymal stem cells post slow-freezing or vitrification. <u>Acta Histochem. 121 (4): 491-9.</u> Desando, G. <i>et al.</i> (2021) Prospects on the Potential In Vitro Regenerative Features of Mechanically Treated-Adipose Tissue for Osteoarthritis Care. <u>Stem Cell Rev Rep. 17 (4):</u>
	<u>1362-73.</u> 21. Adly, H. <i>et al.</i> (2021) Repopulation of multipotent stem cells derived from adult male rabbits on a polycaprolactone scaffold: an <i>in vitro.</i> study <u>Al-Azhar International Medical</u> Journal. 2 (11): 37-42.
	 22. Wee, J.H. <i>et al.</i> (2022) Stem cell laden nano and micro collagen/PLGA bimodal fibrous patches for myocardial regeneration. <u>Biomater Res. 26 (1): 79.</u> 23. Mastrolia, I. <i>et al.</i> (2022) Autologous Marrow Mesenchymal Stem Cell Driving Bone Regeneration in a Rabbit Model of Femoral Head Osteonecrosis. <u>Pharmaceutics.</u> 14(10):2127.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA808GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)RPEGoat Anti Mouse IgG (H/L) (STAR117...)FITCRabbit Anti Mouse IgG (STAR13...)HRPRabbit Anti Mouse IgG (STAR9...)FITC

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rad	.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381507:210512'

Printed on 06 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint