

Datasheet: MCA808GA

BATCH NUMBER 164282

Description:	MOUSE ANTI RABBIT CD45
Specificity:	CD45
Other names:	LCA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	L12/201
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0mg/ml

Immunogen Glycoproteins isolated from the T cell line, RL-5.

RRID AB_10961760

Fusion Partners Spleen cells from immunised mice were fused with cells of the P3.X63.Ag8-U1 mouse myeloma cell line.

Specificity **Mouse anti Rabbit CD45 antibody, clone L12/201** recognizes the CD45 antigen, also known as leukocyte common antigen (LCA) or T200. Mouse anti Rabbit CD45 antibody, clone L12/201 shows pan leucocyte reactivity by flow cytometry and immunohistochemistry.

Immunoprecipitation was achieved by cross linking antibody to the labelled cell surface yielding a protein migrating by gel electrophoresis at a molecular mass of ~200 kDa.

Flow Cytometry Use 10ul of the suggested working dilution to label 10^6 cells or cells or 100ul whole blood.

References

1. Jackson, S. *et al.* (1983) Differentiation antigens identify subpopulations of rabbit T and B lymphocytes. Definition by flow cytometry. [J Exp Med. 157 \(1\): 34-46.](#)
2. Wilkinson, J.M. *et al.* (1984) Cell surface glycoproteins of rabbit lymphocytes: characterization with monoclonal antibodies. [Mol Immunol. 21 \(1\): 95-103.](#)
3. Wilkinson, J.M. *et al.* (1992) A cytotoxic rabbit T-cell line infected with a gamma-herpes virus which expresses CD8 and class II antigens. [Immunology. 77 \(1\): 106-8.](#)
4. Wilkinson, J.M. *et al.* (1993) Immunohistochemical identification of leucocyte populations in normal tissue and inflamed synovium of the rabbit. [J Pathol. 170 \(3\): 315-20.](#)
5. Xu, Y. *et al.* (2010) Adenovirus-mediated overexpression of glutathione-s-transferase mitigates transplant arteriosclerosis in rabbit carotid allografts. [Transplantation. 89: 409-16.](#)
6. Mackenzie, S.M. *et al.* (2006) Immunocontraceptive effects on female rabbits infected with recombinant myxoma virus expressing rabbit ZP2 or ZP3. [Biol Reprod. 74: 511-21.](#)
7. Fenton, M. *et al.* (2001) Cellular senescence after single and repeated balloon catheter denudations of rabbit carotid arteries. [Arterioscler Thromb Vasc Biol. 21: 220-6.](#)
8. Liang, H. *et al.* (2009) Comparison of the ocular tolerability of a latanoprost cationic emulsion versus conventional formulations of prostaglandins: an *in vivo* toxicity assay. [Mol Vis. 15: 1690-9.](#)
9. Wacławicek, M. *et al.* (2009) Analysis of the early response to TSST-1 reveals Vbeta-unrestricted extravasation, compartmentalization of the response, and unresponsiveness but not energy to TSST-1. [J Leukoc Biol. 85: 44-54.](#)
10. Kuznetsov, S.A. *et al.* (2001) Circulating skeletal stem cells. [J Cell Biol. 153: 1133-40.](#)
11. Sijnave, D. *et al.* (2015) Inhibition of Rho-Associated Kinase Prevents Pathological

- Wound Healing and Neovascularization After Corneal Trauma. [Cornea. 34 \(9\): 1120-9.](#)
12. Kováč, M. *et al.* (2016) Cryopreservation of Amniotic Fluid Stem Cells Derived From Zobor Rabbits. [Slovak J Anim Sci., 49,\(2\): 62–67.](#)
13. Ondruska, L.. *et al.* (2016) Decrease in C-reactive protein levels in rabbits after vaccination with a live attenuated myxoma virus vaccine [Veterinárni Medicína. 61 \(No. 10\): 571-6.](#)
14. Davis, W.C. & Hamilton, M.J. (2008) Use of flow cytometry to develop and characterize a set of monoclonal antibodies specific for rabbit leukocyte differentiation molecules. [J Vet Sci. 9 \(1\): 51-66.](#)
15. Waclavicek, M. *et al.* (2009) Analysis of the early response to TSST-1 reveals Vbeta-unrestricted extravasation, compartmentalization of the response, and unresponsiveness but not anergy to TSST-1. [J Leukoc Biol. 85 \(1\): 44-54.](#)
16. Miyahara, T. *et al.* (2013) D-series resolvins attenuates vascular smooth muscle cell activation and neointimal hyperplasia following vascular injury. [FASEB J. 27 \(6\): 2220-32.](#)
17. Vasicek, J. *et al.* (2015) Determination of Lymphocyte Subset Distribution in the Peripheral Blood of Rabbits Immunized with CFA. [Int Symp Anim Sci UDC:639.112 pp. 226-31.](#)
18. Kováč, M. *et al.* (2017) Phenotype and ultrastructure of stem cells derived from amniotic fluid of Nitra rabbit [J Cent Eur Agric.18 \(1\): 226-34.](#)
19. Vašíček, J *et al.* (2018) The Efficiency of Immunomagnetic Sorting of Rabbit Bone Marrow Cells for the Establishment of Mesenchymal Stem Cell Culture. [J Microbiol Biotech Food Sci. 8 \(3\): 890-2.](#)
20. Vašíček, J *et al.* (2014) Basic Blood Analysis of Rabbits Immunized with Vaccine Against Myxomatosis. [Proc Int Symp Anim Sci 2014: 411-6.](#)
21. Ondruska, L. *et al.* (2016) Decrease in C-reactive protein levels in rabbits after vaccination with a live attenuated myxoma virus vaccine [Veterinárni Medicína. 61 \(No. 10\): 571-6.](#)
22. Barth, H. *et al.* (2019) Inflammatory responses after vitrectomy with vitreous substitutes in a rabbit model. [Graefes Arch Clin Exp Ophthalmol. 257 \(4\): 769-83.](#)
23. Kovac, M. *et al.* (2017) Different RNA and protein expression of surface markers in rabbit amniotic fluid-derived mesenchymal stem cells. [Biotechnol Prog. 33 \(6\): 1601-13.](#)
24. Desando, G. *et al.* (2021) Prospects on the Potential In Vitro Regenerative Features of Mechanically Treated-Adipose Tissue for Osteoarthritis Care. [Stem Cell Rev Rep. 17 \(4\): 1362-73.](#)
25. Kulikova, B. *et al.* (2019) Survivability of rabbit amniotic fluid-derived mesenchymal stem cells post slow-freezing or vitrification. [Acta Histochem. 121 \(4\): 491-9.](#)
26. Adly, H. *et al.* (2021) Repopulation of multipotent stem cells derived from adult male rabbits on a polycaprolactone scaffold: an *in vitro*. study [Al-Azhar International Medical Journal. 2 \(11\): 37-42.](#)
27. Wee, J.H. *et al.* (2022) Stem cell laden nano and micro collagen/PLGA bimodal fibrous patches for myocardial regeneration. [Biomater Res. 26 \(1\): 79.](#)
28. Mastrolia, I. *et al.* (2022) Autologous Marrow Mesenchymal Stem Cell Driving Bone Regeneration in a Rabbit Model of Femoral Head Osteonecrosis. [Pharmaceutics. 14\(10\):2127.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA808GA 10040
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [FITC](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M381507:210512'

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)