

Datasheet: MCA80A647

Description:	MOUSE ANTI HUMAN CD1a:Alexa Fluor® 647
Specificity:	CD1a
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	NA1/34-HLK
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Dog, Cynomolgus monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Alexa Fluor® 647 - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
Alexa Fluor®647	650	665

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.

Buffer Solution

Phosphate buffered saline

Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml
Immunogen	Human thymocytes
External Database Links	<p>UniProt: P06126 Related reagents</p> <p>Entrez Gene: 909 CD1A Related reagents</p>
RRID	AB_324820
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NS1/1 Ag4.1 mouse myeloma cell line
Specificity	Mouse anti Human CD1a antibody, clone NA1/34-HLK recognizes the human CD1a cell surface glycoprotein, a ~49 kDa single pass type 1 transmembrane glycoprotein containing a single Ig-like domain, expressed in association with beta 2 microglobulin. CD1a is expressed strongly by cortical thymocytes, and also by Langerhans cells and interdigitating cells. CD1a is involved in the presentation of lipids and glycolipids to NK cells (Sloma <i>et al.</i> 2008).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> McMichael, A.J. <i>et al.</i> (1979) A human thymocyte antigen defined by a hybrid myeloma monoclonal antibody. Eur J Immunol. 9 (3): 205-10. Poulter, L.W. <i>et al.</i> (1986) Discrimination of human macrophages and dendritic cells by means of monoclonal antibodies. Scand J Immunol. 24 (3): 351-7. Galkowska, H. <i>et al.</i> (1996) Reactivity of antibodies directed against human antigens with surface markers on canine leukocytes. Vet Immunol Immunopathol. 53 (3-4): 329-34. Yoshino, N. <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. Exp Anim. 49 (2): 97-110. Hirbod, T. <i>et al.</i> (2010) Abundant expression of HIV target cells and C-type lectin receptors in the foreskin tissue of young Kenyan men. Am J Pathol. 176: 2798-805. Liu, C.C. <i>et al.</i> (2008) Transient downregulation of monocyte-derived dendritic-cell differentiation, function, and survival during tumoral progression and regression in an <i>in vivo</i> canine model of transmissible venereal tumor. Cancer Immunol Immunother. 57: 479-91. Sugiura K <i>et al.</i> (2010) Effect of IL-12 on canine dendritic cell maturation following differentiation induced by granulocyte-macrophage CSF and IL-4. Vet Immunol Immunopathol. 137 (3-4): 322-6. Angel, C.E. <i>et al.</i> (2006) Distinctive localization of antigen-presenting cells in human

- lymph nodes. [Blood. 113: 1257-67.](#)
9. Bosco, M.C. *et al.* (2011) Hypoxia modulates the gene expression profile of immunoregulatory receptors in human mature dendritic cells: identification of TREM-1 as a novel hypoxic marker in vitro and *in vivo*. [Blood. 117: 2625-39.](#)
10. Buettner, M. *et al.* (2005) Inverse correlation of maturity and antibacterial activity in human dendritic cells. [J Immunol. 174: 4203-9.](#)
11. Cox, K. *et al.* (2005) Plasmacytoid dendritic cells (PDC) are the major DC subset innately producing cytokines in human lymph nodes. [J Leukoc Biol. 78: 1142-52.](#)
12. Mito, K. *et al.* (2010) IFN γ markedly cooperates with intratumoral dendritic cell vaccine in dog tumor models. [Cancer Res. 70: 7093-101.](#)
13. Murray, S. *et al.* (2000) Diagnostic and therapeutic evaluation of an anti-Langerhans cell histiocytosis monoclonal antibody (NA1/34) in a new xenograft model. [J Invest Dermatol. 114: 127-34.](#)
14. Scheinecker, C. *et al.* (1998) Initiation of the autologous mixed lymphocyte reaction requires the expression of costimulatory molecules B7-1 and B7-2 on human peripheral blood dendritic cells. [J Immunol. 161: 3966-73.](#)
15. Wang, Y.S. *et al.* (2007) Characterization of canine monocyte-derived dendritic cells with phenotypic and functional differentiation. [Can J Vet Res. 71: 165-74.](#)
16. Elia, A.R. *et al.* (2008) Human dendritic cells differentiated in hypoxia down-modulate antigen uptake and change their chemokine expression profile. [J Leukoc Biol. 84: 1472-82.](#)
17. Fanales-Belasio, E. *et al.* (2009) HIV-1 Tat addresses dendritic cells to induce a predominant Th1-type adaptive immune response that appears prevalent in the asymptomatic stage of infection. [J Immunol. 182: 2888-97.](#)
18. Kaldensjö, T. *et al.* (2011) Detection of intraepithelial and stromal Langerin and CCR5 positive cells in the human endometrium: potential targets for HIV infection. [PLoS One. 6: e21344.](#)
19. Angel, C.E. *et al.* (2007) CD14⁺ antigen-presenting cells in human dermis are less mature than their CD1a⁺ counterparts. [Int Immunol. 19: 1271-9.](#)
20. Angel, C.E. *et al.* (2007) Comprehensive analysis of MHC-II expression in healthy human skin. [Immunol Cell Biol. 85: 363-9.](#)
21. Baharom F. *et al.* (2016) Dendritic Cells and Monocytes with Distinct Inflammatory Responses Reside in Lung Mucosa of Healthy Humans [The Journal of Immunology. May 2 \[Epub ahead of print\]](#)
22. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. [Immunobiology. 221 \(1\): 12-22.](#)
23. Zegarska, B. *et al.* (2017) Changes of Langerhans cells during skin ageing. [Postepy Dermatol Alergol. 34 \(3\): 260-7.](#)
24. Tomić, S. *et al.* (2018) Functionalization-dependent effects of cellulose nanofibrils on tolerogenic mechanisms of human dendritic cells. [Int J Nanomedicine. 13: 6941-60.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and

should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA929A647\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M368996:200529'

Printed on 02 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)