

Datasheet: MCA81

BATCH NUMBER 160947

Description:	MOUSE ANTI HUMAN HLA ABC
Specificity:	HLA ABC
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	W6/32
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Macaque, Bovine, Cynomolgus monkey, Baboon, Rhesus Monkey, Chimpanzee, Gorilla, Shrew</p> <p>Does not react with: Goat, Dog, Guinea Pig, Rabbit, Mouse, Chicken, Amphibia</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or</p>

personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified human tonsil lymphocyte membranes.
RRID	AB_322094
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1/1-Ag4.1 myeloma cell line.
Specificity	<p>Mouse anti Human HLA ABC antibody, clone W6/32 recognizes an antigenic determinant shared among products of the HLA A, B and C loci. Clone W6/32 recognizes a conformational epitope, reacting with HLA class I alpha3 and alpha2 domains. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In humans, this complex is referred to as the human leukocyte antigen (HLA) region. There are 3 major MHC class I proteins encoded by the HLA which are HLA A, HLA B and HLA C. These proteins are found on the surface of almost all nucleated somatic cells.</p> <p>Mouse anti Human HLA ABC antibody, clone W6/32 is routinely tested in flow cytometry on human peripheral blood lymphocytes.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells or 100ul whole blood
Histology Positive Control Tissue	Tonsil
References	<ol style="list-style-type: none"> 1. Barnstable, C.J. <i>et al.</i> (1978) Production of monoclonal antibodies to group A erythrocytes, HLA and other human cell surface antigens-new tools for genetic analysis. Cell. 14 (1): 9-20. 2. Jacobsen, C.N. <i>et al.</i> (1993) Reactivities of 20 anti-human monoclonal antibodies with leucocytes from ten different animal species. Vet Immunol Immunopathol. 39 (4): 461-6. 3. Yoshino, N. <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by

- using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)
4. Neefjes, J.J. *et al.* (1986) A biochemical characterization of feline MHC products: unusually high expression of class II antigens on peripheral blood lymphocytes. [Immunogenetics. 23 \(5\): 341-7.](#)
5. Stern, P.L. *et al.* (1987) Class I-like MHC molecules expressed by baboon placental syncytiotrophoblast. [J Immunol. 138 \(4\): 1088-91.](#)
6. Verbeek, M.M. *et al.* (1995) T lymphocyte adhesion to human brain pericytes is mediated via very late antigen-4/vascular cell adhesion molecule-1 interactions. [J Immunol. 154 \(11\): 5876-84.](#)
7. Tanabe, M. *et al.* (1992) Structural and functional analysis of monomorphic determinants recognized by monoclonal antibodies reacting with the HLA class I alpha 3 domain. [J Immunol. 148 \(10\): 3202-9.](#)
8. Ishitani, A. *et al.* (2003) Protein expression and peptide binding suggest unique and interacting functional roles for HLA-E, F, and G in maternal-placental immune recognition. [J Immunol. 171 \(3\): 1376-84.](#)
9. Dressel, R. *et al.* (2003) Differential effect of acute and permanent heat shock protein 70 overexpression in tumor cells on lysability by cytotoxic T lymphocytes. [Cancer Res. 63 \(23\): 8212-20.](#)
10. Brodsky, F.M. & Parham, P. (1982) Evolution of HLA antigenic determinants: species cross-reactions of monoclonal antibodies. [Immunogenetics. 15 \(2\): 151-66.](#)
11. Hinrichs, J. *et al.* (2010) The nature of peptides presented by an HLA class I low expression allele. [Haematologica. 95: 1373-80.](#)
12. Jones, D.C. *et al.* (2011) HLA Class I Allelic Sequence and Conformation Regulate Leukocyte Ig-Like Receptor Binding. [J Immunol. 186: 2990-7.](#)
13. Spentzou, A. *et al.* (2010) Viral inhibition assay: a CD8 T cell neutralization assay for use in clinical trials of HIV-1 vaccine candidates. [J Infect Dis. 201: 720-9.](#)
14. Fujita, Y. *et al.* (2010) Bone marrow transplantation restores epidermal basement membrane protein expression and rescues epidermolysis bullosa model mice. [Proc Natl Acad Sci U S A. 107: 14345-50.](#)
15. Grotzke, J.E. *et al.* (2009) The Mycobacterium tuberculosis phagosome is a HLA-I processing competent organelle. [PLoS Pathog. 5: e1000374.](#)
16. Narita, M. *et al.* (2010) WT1 peptide vaccination in combination with imatinib therapy for a patient with CML in the chronic phase. [Int J Med Sci. 7: 72-81.](#)
17. Vitadello, M. *et al.* (2010) Myofiber stress-response in myositis: parallel investigations on patients and experimental animal models of muscle regeneration and systemic inflammation. [Arthritis Res Ther. 12: R52.](#)
18. Zuo, J. *et al.* (2011) The Epstein-Barr virus-encoded BILF1 protein modulates immune recognition of endogenously processed antigen by targeting major histocompatibility complex class I molecules trafficking on both the exocytic and endocytic pathways. [J Virol. 85: 1604-14.](#)
19. Enose-Akahata, Y. *et al.* (2012) Minocycline modulates antigen-specific CTL activity through inactivation of mononuclear phagocytes in patients with HTLV-I associated neurologic disease. [Retrovirology. 9: 16.](#)
20. Badrinath, S. *et al.* (2012) Position 156 influences the peptide repertoire and tapasin dependency of human leukocyte antigen B*44 allotypes. [Haematologica. 97: 98-106.](#)
21. Tischer, S. *et al.* (2016) Discovery of immunodominant T-cell epitopes reveals penton protein as a second immunodominant target in human adenovirus infection. [J Transl Med.](#)

[14 \(1\): 286.](#)

22. Dragovic, R.A. *et al.* (2015) Isolation of syncytiotrophoblast microvesicles and exosomes and their characterisation by multicolour flow cytometry and fluorescence Nanoparticle Tracking Analysis. [Methods. 87: 64-74.](#)

23. Praest, P. *et al.* (2019) A Flow Cytometry-Based Approach to Unravel Viral Interference with the MHC Class I Antigen Processing and Presentation Pathway. [Methods Mol Biol. 1988: 187-98.](#)

24. Tannetta, D.S. *et al.* (2013) Characterisation of syncytiotrophoblast vesicles in normal pregnancy and pre-eclampsia: expression of Flt-1 and endoglin. [PLoS One. 8 \(2\): e56754.](#)

25. Juan, C.H. *et al.* (2020) In Vitro Differentiation of Human Placenta-Derived Multipotent Cells into Schwann-Like Cells. [Biomolecules. 10 \(12\): 1657.](#)

26. Tupova, L. *et al.* (2020) Interplay of drug transporters P-glycoprotein (MDR1), MRP1, OATP1A2 and OATP1B3 in passage of maraviroc across human placenta. [Biomed Pharmacother. 129: 110506.](#)

27. Nguyen, J. *et al.* (2021) Quantitative contributions of TNF receptor superfamily members to CD8⁺ T-cell responses. [Mol Syst Biol. 17 \(11\): e10560.](#)

28. Cacciola, R. *et al.* (2022) Impact of Anti-Endothelial Cell Antibodies (AECAs) in Patients with Polycythemia Vera and Thrombosis [Diagnostics. 12 \(5\): 1077.](#)

29. Xia, P. *et al.* (2022) NLRP3 inflammasome up-regulates major histocompatibility complex class I expression and promotes inflammatory infiltration in polymyositis. [BMC Immunol. 23 \(1\): 39.](#)

30. Korbonits, L. *et al.* (2022) *Mycobacterium avium* subsp. *paratuberculosis* Infected Cows Reveal Divergent Immune Response in Bovine Peripheral Blood Derived Lymphocyte Proteome. [Metabolites. 12 \(10\): 924.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA81>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M382566:210513'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)