

Datasheet: MCA81PE

Description:	MOUSE ANTI HUMAN HLA ABC:RPE
Specificity:	HLA ABC
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	W6/32
Isotype:	lgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Macaqu	e, Bovine, Cynomolgus	s monkey, Baboon,	Rhesus Monkey,
Reactivity	Chimpanzee, Gorilla,	, Shrew		
	Does not react with:	Goat, Dog, Guinea Pig,	Rabbit, Mouse, Ch	icken, Amphibia
	-	vity and working conditi		
	•	rom testing within our la tions from the originato	• •	•
Product Form	personal communica further information.	J	ors. Please refer to	references indicated
Product Form Reconstitution	personal communica further information.	tions from the originato	ors. Please refer to	references indicated
	personal communica further information. Purified IgG conjugat	tions from the originato	ors. Please refer to	references indicated

supernatant.

·
Phosphate buffered saline
0.09% Sodium Azide
1% Bovine Serum Albumin
5% Sucrose
Purified human tonsil lymphocyte membranes.
AB_322096
Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1/1-Ag4.1 myeloma cell line.
Mouse anti Human HLA ABC antibody, clone W6/32 recognizes an antigenic determinant shared among products of the HLA A, B and C loci. Clone W6/32 recognizes a conformational epitope, reacting with HLA class I alpha3 and alpha2 domains. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In humans, this complex is referred to as the human leukocyte antigen (HLA) region. There are 3 major MHC class I proteins encoded by the HLA which are HLA A, HLA B and HLA C. These proteins are found on the surface of almost all nucleated somatic cells.
Mouse anti Human HLA ABC antibody, clone W6/32 is routinely tested in flow cytometry on human peripheral blood lymphocytes.
Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
 Brodsky, F.M. & Parham, P. (1982) Evolution of HLA antigenic determinants: species cross-reactions of monoclonal antibodies. <u>Immunogenetics</u>. 15 (2): 151-66. Neefjes, J.J. <i>et al.</i> (1986) A biochemical characterization of feline MHC products: unusually high expression of class II antigens on peripheral blood lymphocytes. <u>Immunogenetics</u>. 23 (5): 341-7. Stern, P.L. <i>et al.</i> (1987) Class I-like MHC molecules expressed by baboon placental syncytiotrophoblast. <u>J Immunol</u>. 138 (4): 1088-91. Jacobsen, C.N. <i>et al.</i> (1993) Reactivities of 20 anti-human monoclonal antibodies with leucocytes from ten different animal species. <u>Vet Immunol Immunopathol</u>. 39 (4): 461-6. Verbeek, M.M. <i>et al.</i> (1995) T lymphocyte adhesion to human brain pericytes is mediated via very late antigen-4/vascular cell adhesion molecule-1 interactions. <u>J</u>

8. Ishitani, A. *et al.* (2003) Protein expression and peptide binding suggest unique and interacting functional roles for HLA-E, F, and G in maternal-placental immune recognition. <u>J Immunol. 171 (3): 1376-84.</u>

- 9. Giuliani, F. *et al.* (2003) Vulnerability of human neurons to T cell-mediated cytotoxicity. <u>J</u> Immunol. 171: 368-79.
- 10. Smith, A.C. *et al.* (2005) Interaction of the *Salmonella*-containing vacuole with the endocytic recycling system. <u>J Biol Chem. 280: 24634-41.</u>
- 11. Zuo, J. *et al.* (2008) The DNase of gammaherpesviruses impairs recognition by virus-specific CD8+ T cells through an additional host shutoff function. J Virol. 82: 2385-93.
- 12. Shi, J. *et al.* (2008) Bortezomib down-regulates the cell-surface expression of HLA class I and enhances natural killer cell-mediated lysis of myeloma. <u>Blood. 111: 1309-17.</u>
- 13. Watling, D. *et al.* (2008) Multiple kinases in the interferon-gamma response. <u>Proc Natl</u> Acad Sci U S A. 105: 6051-6.
- 14. Grotzke, J.E. *et al.* (2009) The *Mycobacterium tuberculosis* phagosome is a HLA-I processing competent organelle. PLoS Pathog. 5: e1000374.
- 15. Koch, N. *et al.* (2009) IL-10 protects monocytes and macrophages from complement-mediated lysis. J Leukoc Biol. 86: 155-66.
- 16. Spentzou, A. *et al.* (2010) Viral inhibition assay: a CD8 T cell neutralization assay for use in clinical trials of HIV-1 vaccine candidates. J Infect Dis. 201: 720-9.
- 17. Vitadello, M. *et al.* (2010) Myofiber stress-response in myositis: parallel investigations on patients and experimental animal models of muscle regeneration and systemic inflammation. Arthritis Res Ther. 12: R52.
- 18. Hinrichs, J. *et al.* (2010) The nature of peptides presented by an HLA class I low expression allele. <u>Haematologica</u>. 95: 1373-80.
- 19. Fujita, Y. *et al.* (2010) Bone marrow transplantation restores epidermal basement membrane protein expression and rescues epidermolysis bullosa model mice. <u>Proc Natl Acad Sci U S A. 107: 14345-50.</u>
- 20. Narita, M. *et al.* (2010) WT1 peptide vaccination in combination with imatinib therapy for a patient with CML in the chronic phase. <u>Int J Med Sci. 7: 72-81.</u>
- 21. Zuo, J. *et al.* (2011) The Epstein-Barr virus-encoded BILF1 protein modulates immune recognition of endogenously processed antigen by targeting major histocompatibility complex class I molecules trafficking on both the exocytic and endocytic pathways. <u>J Virol.</u> 85: 1604-14.
- 22. Lask, A. *et al.* (2011) TCR-Independent Killing of B Cell Malignancies by Anti-Third-Party CTLs: The Critical Role of MHC-CD8 Engagement. <u>J Immunol. 187:</u> 2006-14.
- 23. Jones, D.C. *et al.* (2011) HLA Class I Allelic Sequence and Conformation Regulate Leukocyte Ig-Like Receptor Binding. <u>J Immunol</u>. 186: 2990-7.
- 24. Enose-Akahata, Y. *et al.* (2012) Minocycline modulates antigen-specific CTL activity through inactivation of mononuclear phagocytes in patients with HTLV-I associated neurologic disease. <u>Retrovirology. 9: 16.</u>
- 25. Badrinath, S. *et al.* (2012) Position 156 influences the peptide repertoire and tapasin dependency of human leukocyte antigen B*44 allotypes. <u>Haematologica</u>. 97: 98-106.
- 26. Tannetta, D.S. *et al.* (2013) Characterisation of syncytiotrophoblast vesicles in normal pregnancy and pre-eclampsia: expression of Flt-1 and endoglin. <u>PLoS One. 8 (2): e56754.</u>
- 27. Dragovic, R.A. *et al.* (2015) Isolation of syncytiotrophoblast microvesicles and exosomes and their characterisation by multicolour flow cytometry and fluorescence Nanoparticle Tracking Analysis. <u>Methods</u>. 87: 64-74.
- 28. Tischer, S. *et al.* (2016) Discovery of immunodominant T-cell epitopes reveals penton protein as a second immunodominant target in human adenovirus infection. <u>J Transl Med.</u>

14 (1): 286.

- 29. Praest, P. *et al.* (2019) A Flow Cytometry-Based Approach to Unravel Viral Interference with the MHC Class I Antigen Processing and Presentation Pathway. Methods Mol Biol. 1988: 187-98.
- 30. Juan, C.H. *et al.* (2020) *In Vitro* Differentiation of Human Placenta-Derived Multipotent Cells into Schwann-Like Cells. Biomolecules. 10 (12): 1657.
- 31. Tupova, L. *et al.* (2020) Interplay of drug transporters P-glycoprotein (MDR1), MRP1, OATP1A2 and OATP1B3 in passage of maraviroc across human placenta. <u>Biomed Pharmacother</u>. 129: 110506.
- 32. Nguyen, J. *et al.* (2021) Quantitative contributions of TNF receptor superfamily members to CD8⁺ T-cell responses. Mol Syst Biol. 17 (11): e10560.
- 33. Xia, P. *et al.* (2022) NLRP3 inflammasome up-regulates major histocompatibility complex class I expression and promotes inflammatory infiltration in polymyositis. <u>BMC Immunol. 23 (1): 39.</u>
- 34. Cacciola, R. *et al.* (2022) Impact of Anti-Endothelial Cell Antibodies (AECAs) in Patients with Polycythemia Vera and Thrombosis Diagnostics. 12 (5): 1077.
- 35. Korbonits, L. *et al.* (2022) *Mycobacterium avium* subsp. *paratuberculosi*s Infected Cows Reveal Divergent Immune Response in Bovine Peripheral Blood Derived Lymphocyte Proteome. <u>Metabolites.12 (10): 924.</u>
- 36. Lyssy, F. *et al.* (2023) Platelet-derived factors dysregulate placental sphingosine-1-phosphate receptor 2 in human trophoblasts. Reprod Biomed Online. 47 (2): 103215.
- 37. Forstner, D. *et al.* (2023) CD39 abrogates platelet-derived factors induced IL-1 β expression in the human placenta. <u>Front Cell Dev Biol. 11: 1183793.</u>
- 38. Zheng, S. *et al.* (2022) The SARS-CoV-2 accessory factor ORF7a downregulates MHC class I surface expression. <u>bioRxiv 30 May [Epub ahead of print. Preprint]</u>
- 39. Fielding, C.A. *et al.* (2022) SARS-CoV-2 host-shutoff impacts innate NK cell functions, but antibody-dependent NK activity is strongly activated through non-spike antibodies. <u>11:</u> <u>e74489.</u>
- 40. Santamorena, M.M. *et al.* (2024) Engineered HCMV-infected APCs enable the identification of new immunodominant HLA-restricted epitopes of anti-HCMV T-cell immunity. <u>HLA. 103 (6): e15541.</u>

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA81PE 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:RPE (MCA929PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

 $\textbf{Email: antibody_sales_us@bio-rad.com} \\ \textbf{Email: antibody_sales_uk@bio-rad.com} \\ \textbf{Email: antibody_sales_uk@b$

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419692:230616'

Printed on 14 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint