

Datasheet: MCA81SBR815

## **BATCH NUMBER 64591943**

| Description:  | MOUSE ANTI HUMAN HLA ABC:StarBright Red 815 |
|---------------|---|
| Specificity:  | HLA ABC                                     |
| Format:       | StarBright Red 815                          |
| Product Type: | Monoclonal Antibody                         |
| Clone:        | W6/32                                       |
| Isotype:      | IgG2a                                       |
| Quantity:     | 100 TESTS/0.5ml                             |
|               |   |

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry |     |    |                | Neat               |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species              | Human  |   |   |                     |
|-----------------------------|--|---|---|---------------------|
| Species Cross<br>Reactivity | Reacts with: Macaque, Bovine, Cynomolgus monkey, Baboon, Rhesus Monkey, Chimpanzee, Gorilla, Shrew Does not react with:Goat, Dog, Guinea Pig, Rabbit, Mouse, Chicken, Amphibia |   |   |                     |
|                             | <b>N.B.</b> Antibody reactive reactivity is derived f  | vity and working condit<br>rom testing within our I | ions may vary betwo<br>aboratories, peer-re | •                   |
| Product Form                | Purified IgG conjuga   | ted to StarBright Red 8                             | 315 - liquid                                |                     |
| Max Ex/Em                   | Fluorophore  | Excitation Max (nm)                                 | Emission Max (nm                            | )                   |
|                             | StarBright Red 815   | 654   | 811   |                     |
| Preparation                 | Purified IgG prepare supernatant   | d by affinity chromatog                             | raphy on Protein G                          | from tissue culture |

| Buffer Solution | Phosphate buffered saline   |
|-----------------|---|
| Preservative    | 0.09% Sodium Azide (NaN <sub>3</sub> )  |
| Stabilisers     | 1% Bovine Serum Albumin   |
|                 | 0.1% Pluronic F68   |
|                 | 0.1% PEG 3350   |
|                 | 0.05% Tween 20  |
| Immunogen       | Purified human tonsil lymphocyte membranes.   |
| Fusion Partners | Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1/1-Ag4.1 myeloma cell line.   |
| Specificity     | Mouse anti Human HLA ABC antibody, clone W6/32 recognizes an antigenic determinant shared among products of the HLA A, B and C loci. Clone W6/32 recognizes a conformational epitope, reacting with HLA class I alpha3 and alpha2 domains. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In humans, this complex is referred to as the human leukocyte antigen (HLA) region. There are 3 major MHC class I proteins encoded by the HLA which are HLA A, HLA B and HLA C. These proteins are found on the surface of almost all nucleated somatic cells. |
|                 | Mouse anti Human HLA ABC antibody, clone W6/32 is routinely tested in flow cytometry on human peripheral blood lymphocytes.   |
| Flow Cytometry  | Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.   |
| References      | 1. Brodsky, F.M. & Parham, P. (1982) Evolution of HLA antigenic determinants: species   |
|                 | cross-reactions of monoclonal antibodies. <u>Immunogenetics. 15 (2): 151-66.</u>  |
|                 | 2. Neefjes, J.J. et al. (1986) A biochemical characterization of feline MHC products:   |
|                 | unusually high expression of class II antigens on peripheral blood lymphocytes.   |
|                 | Immunogenetics. 23 (5): 341-7.  |
|                 | 3. Stern, P.L. et al. (1987) Class I-like MHC molecules expressed by baboon placental   |
|                 | syncytiotrophoblast. <u>J Immunol. 138 (4): 1088-91.</u>  |
|                 | 4. Jacobsen, C.N. et al. (1993) Reactivities of 20 anti-human monoclonal antibodies with  |
|                 | leucocytes from ten different animal species. <u>Vet Immunol Immunopathol. 39 (4): 461-6.</u>   |
|                 | 5. Verbeek, M.M. et al. (1995) T lymphocyte adhesion to human brain pericytes is  |
|                 | mediated via very late antigen-4/vascular cell adhesion molecule-1 interactions. <u>J</u>   |
|                 | <u>Immunol. 154 (11): 5876-84.</u>  |
|                 | 6. Raftery, M.J. et al. (2002) Hantavirus infection of dendritic cells. <u>J Virol. 76: 10724-33.</u>   |
|                 | 7. Dressel, R. et al. (2003) Differential effect of acute and permanent heat shock protein  |
|                 | 70 overexpression in tumor cells on lysability by cytotoxic T lymphocytes. <u>Cancer Res. 63</u>  |
|                 | (23): 8212-20   |

- (23): 8212-20.
- 8. Ishitani, A. et al. (2003) Protein expression and peptide binding suggest unique and interacting functional roles for HLA-E, F, and G in maternal-placental immune recognition. J Immunol. 171 (3): 1376-84.
- 9. Giuliani, F. et al. (2003) Vulnerability of human neurons to T cell-mediated cytotoxicity. J

#### Immunol. 171: 368-79.

- 10. Smith, A.C. *et al.* (2005) Interaction of the *Salmonella*-containing vacuole with the endocytic recycling system. J Biol Chem. 280: 24634-41.
- 11. Zuo, J. *et al.* (2008) The DNase of gammaherpesviruses impairs recognition by virus-specific CD8+ T cells through an additional host shutoff function. <u>J Virol. 82: 2385-93.</u>
- 12. Shi, J. *et al.* (2008) Bortezomib down-regulates the cell-surface expression of HLA class I and enhances natural killer cell-mediated lysis of myeloma. Blood. 111: 1309-17.
- 13. Watling, D. *et al.* (2008) Multiple kinases in the interferon-gamma response. <u>Proc Natl Acad Sci U S A. 105: 6051-6.</u>
- 14. Grotzke, J.E. *et al.* (2009) The *Mycobacterium tuberculosis* phagosome is a HLA-I processing competent organelle. PLoS Pathog. 5: e1000374.
- 15. Koch, N. *et al.* (2009) IL-10 protects monocytes and macrophages from complement-mediated lysis. J Leukoc Biol. 86: 155-66.
- 16. Spentzou, A. *et al.* (2010) Viral inhibition assay: a CD8 T cell neutralization assay for use in clinical trials of HIV-1 vaccine candidates. J Infect Dis. 201: 720-9.
- 17. Vitadello, M. *et al.* (2010) Myofiber stress-response in myositis: parallel investigations on patients and experimental animal models of muscle regeneration and systemic inflammation. Arthritis Res Ther. 12: R52.
- 18. Hinrichs, J. *et al.* (2010) The nature of peptides presented by an HLA class I low expression allele. <u>Haematologica</u>. 95: 1373-80.
- 19. Fujita, Y. *et al.* (2010) Bone marrow transplantation restores epidermal basement membrane protein expression and rescues epidermolysis bullosa model mice. <u>Proc Natl Acad Sci U S A. 107: 14345-50.</u>
- 20. Narita, M. *et al.* (2010) WT1 peptide vaccination in combination with imatinib therapy for a patient with CML in the chronic phase. Int J Med Sci. 7: 72-81.
- 21. Zuo, J. *et al.* (2011) The Epstein-Barr virus-encoded BILF1 protein modulates immune recognition of endogenously processed antigen by targeting major histocompatibility complex class I molecules trafficking on both the exocytic and endocytic pathways. <u>J Virol.</u> 85: 1604-14.
- 22. Lask, A. *et al.* (2011) TCR-Independent Killing of B Cell Malignancies by Anti-Third-Party CTLs: The Critical Role of MHC-CD8 Engagement. <u>J Immunol. 187:</u> 2006-14.
- 23. Jones, D.C. *et al.* (2011) HLA Class I Allelic Sequence and Conformation Regulate Leukocyte Ig-Like Receptor Binding. <u>J Immunol</u>. 186: 2990-7.
- 24. Enose-Akahata, Y. *et al.* (2012) Minocycline modulates antigen-specific CTL activity through inactivation of mononuclear phagocytes in patients with HTLV-I associated neurologic disease. <u>Retrovirology. 9: 16.</u>
- 25. Badrinath, S. *et al.* (2012) Position 156 influences the peptide repertoire and tapasin dependency of human leukocyte antigen B\*44 allotypes. <u>Haematologica</u>. 97: 98-106.
- 26. Tannetta, D.S. *et al.* (2013) Characterisation of syncytiotrophoblast vesicles in normal pregnancy and pre-eclampsia: expression of Flt-1 and endoglin. <u>PLoS One. 8 (2): e56754.</u>
- 27. Dragovic, R.A. *et al.* (2015) Isolation of syncytiotrophoblast microvesicles and exosomes and their characterisation by multicolour flow cytometry and fluorescence Nanoparticle Tracking Analysis. <u>Methods. 87: 64-74.</u>
- 28. Tischer, S. *et al.* (2016) Discovery of immunodominant T-cell epitopes reveals penton protein as a second immunodominant target in human adenovirus infection. <u>J Transl Med.</u> 14 (1): 286.

- 29. Praest, P. *et al.* (2019) A Flow Cytometry-Based Approach to Unravel Viral Interference with the MHC Class I Antigen Processing and Presentation Pathway. Methods Mol Biol. 1988: 187-98.
- 30. Juan, C.H. *et al.* (2020) *In Vitro* Differentiation of Human Placenta-Derived Multipotent Cells into Schwann-Like Cells. <u>Biomolecules</u>. 10 (12): 1657.
- 31. Tupova, L. *et al.* (2020) Interplay of drug transporters P-glycoprotein (MDR1), MRP1, OATP1A2 and OATP1B3 in passage of maraviroc across human placenta. <u>Biomed Pharmacother</u>. 129: 110506.
- 32. Nguyen, J. *et al.* (2021) Quantitative contributions of TNF receptor superfamily members to CD8<sup>+</sup> T-cell responses. <u>Mol Syst Biol. 17 (11): e10560.</u>
- 33. Xia, P. *et al.* (2022) NLRP3 inflammasome up-regulates major histocompatibility complex class I expression and promotes inflammatory infiltration in polymyositis. <u>BMC Immunol.</u> 23 (1): 39.
- 34. Cacciola, R. *et al.* (2022) Impact of Anti-Endothelial Cell Antibodies (AECAs) in Patients with Polycythemia Vera and Thrombosis <u>Diagnostics</u>. 12 (5): 1077.
- 35. Korbonits, L. *et al.* (2022) *Mycobacterium avium* subsp. *paratuberculosi*s Infected Cows Reveal Divergent Immune Response in Bovine Peripheral Blood Derived Lymphocyte Proteome. <u>Metabolites.12 (10): 924.</u>
- 36. Lyssy, F. *et al.* (2023) Platelet-derived factors dysregulate placental sphingosine-1-phosphate receptor 2 in human trophoblasts. Reprod Biomed Online. 47 (2): 103215.
- 37. Forstner, D. *et al.* (2023) CD39 abrogates platelet-derived factors induced IL-1β expression in the human placenta. <u>Front Cell Dev Biol.</u> 11: 1183793.
- 38. Zheng, S. *et al.* (2022) The SARS-CoV-2 accessory factor ORF7a downregulates MHC class I surface expression. <u>bioRxiv 30 May [Epub ahead of print. Preprint]</u>
- 39. Fielding, C.A. *et al.* (2022) SARS-CoV-2 host-shutoff impacts innate NK cell functions, but antibody-dependent NK activity is strongly activated through non-spike antibodies. <u>11:</u> e74489.
- 40. Santamorena, M.M. *et al.* (2024) Engineered HCMV-infected APCs enable the identification of new immunodominant HLA-restricted epitopes of anti-HCMV T-cell immunity. <u>HLA. 103 (6): e15541.</u>

| Storage                          | Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.   |
|----------------------------------|--|
| Guarantee                        | 12 months from date of despatch  |
| Acknowledgements                 | This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #20471 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA81SBR815">https://www.bio-rad-antibodies.com/SDS/MCA81SBR815</a> 20471 |
| Regulatory                       | For research purposes only   |

## **Related Products**

#### **Recommended Useful Reagents**

# HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M429010:240308'

## Printed on 12 Dec 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint