

# Datasheet: MCA832F BATCH NUMBER 160952

Description:	MOUSE ANTI BOVINE CD45:FITC			
Specificity:	CD45			
Format:	FITC			
Product Type:	Monoclonal Antibody			
Clone:	CC1			
Isotype:	lgG1			
Quantity:	100 TESTS			

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			Neat - 1/10		
	Where this antibody has not been tested for use in a particular technique this does necessarily exclude its use in such procedures. Suggested working dilutions are g a guide only. It is recommended that the user titrates the antibody for use in their of system using appropriate negative/positive controls.						
Target Species	Bovine						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)			
	FITC	490		525	-		
Preparation	Purified IgG prepared supernatant	by affinity chro	omatogra	aphy on Protein A fro	om tissue culture		
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.1r	ng/ml					

Immunogen	Bovine thoracic duct lymphocytes.				
RRID	AB_2174255				
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.				
Specificity	Mouse anti Bovine CD45 antibody, clone CC1 recognizes the bovine homologue of the human CD45 antigen, also known as leucocyte common antigen (LCA),				
	CD45 is a pan leucocyte cell surface marker expressed on all cells of hematopoietic origin except for erythrocytes ( <u>Bembridge <i>et al.</i> 1993</u> ). CD45 occurs in a number of isoforms, which in bovine includes CD45R, CD45RA, CD45RB and CD45RO and which have restricted cellular expression.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.				
References	<ol> <li>Bembridge, G.P. <i>et al.</i> (1993) Identification of monoclonal antibodies specific for bovine leukocyte common antigen (CD45) together with a novel broadly expressed leukocyte differentiation antigen, BoWC11. <u>Vet Immunol Immunopathol. 39</u>: 115-20.</li> <li>Inchaisri, C. <i>et al.</i> (2000) Studies on the modulation of leucocyte subpopulations and immunoglobulins following intramammary infusion of beta 1,3-glucan into the bovine udder during the dry period. <u>J Vet Med B Infect Dis Vet Public Health. 47 (5)</u>: 373-86.</li> <li>Hu, S. <i>et al.</i> (2001) Effect of subcutaneous injection of ginseng on cows with subclinical <i>Staphylococcus aureus</i> mastitis. <u>J Vet Med B Infect Dis Vet Public Health. 48 (7): 519-28.</u></li> <li>Gånheim, C. <i>et al.</i> (2005) Changes in peripheral blood leucocyte counts and subpopulations after experimental infection with BVDV and/or <i>Mannheimia haemolytica.</i> <u>J Vet Med B Infect Dis Vet Public Health. 52 (9): 380-5.</u></li> <li>Niku, M. <i>et al.</i> (2006) Identification of major cell types in paraffin sections of bovine tissues. <u>BMC Vet Res. 2: 5.</u></li> <li>Grönlund, U. <i>et al.</i> (2008) Effects of road transportation on lymphocyte sub-populations during acute and chronic phases of <i>Staphylococcus aureus</i> induced bovine mastitis. <u>Res Vet Sci. 80 (2): 147-54.</u></li> <li>Riondato, F. <i>et al.</i> (2010) A hierarchy of endothelial colony-forming cell activity displayed by bovine corneal endothelial cells. <u>Invest Ophthalmol Vis Sci. 51: 3943-9.</u></li> <li>Spalenza, V. <i>et al.</i> (2011) Identification of internal control genes for quantitative expression analysis by real-time PCR in bovine peripheral lymphocytes. <u>Vet J. 189 (3): 278-83.</u></li> <li>Lynch, E.M. <i>et al.</i> (2017) Local immunization impacts the response of dairy cows to <i>Escherichia coli</i> mastitis. <u>Sci Rep. 7 (1): 3441.</u></li> <li>Lee, J. <i>et al.</i> (2020) Bovine tongue epithelium-derived cells: A new source of bovine mesenchymal stem cells. <u>Biosci Rep. 40 (4): BSR20181829.</u></li> </ol>				

	13. Molinos, M. <i>et al.</i> (2023) Alterations of bovine nucleus pulposus cells with aging. <u>Aging</u> <u>Cell. 22 (8): e13873.</u>
Further Reading	1. Ballingall, K.T. <i>et al.</i> (2001) The CD45 locus in cattle: allelic polymorphism and evidence for exceptional positive natural selection. <u>Immunogenetics. 52: 276-83.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA832F 10041
Regulatory	For research purposes only

## **Related Products**

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bi	o-rad.com	Email: antibody_sales_uk@bic	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M395822:220519'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint