

Datasheet: MCA837GA

Description:	MOUSE ANTI BOVINE CD8
Specificity:	CD8
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC63
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Species Cross Reactivity	<p>Reacts with: Sheep, Goat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
External Database Links	<p>UniProt: P31783 Related reagents</p> <p>Entrez Gene: 281060 CD8A Related reagents</p>
Fusion Partners	Spleen cells from an immunized mouse were fused with cells of the mouse NS1 myeloma cell line.
Specificity	Mouse anti Bovine CD8 antibody, clone CC63 reacts with the bovine CD8 antigen expressed by a subset of T lymphocytes. The antibody precipitates molecules of ~34 kDa and ~38 kDa under reducing conditions. Clone CC63 has been reported as being suitable for use on formalin dichromate (FD5) fixed paraffin embedded tissue with amplification and antigen retrieval techniques (Gutierrez et al. 1999).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. MacHugh, N.D. & Sopp P (1991) Individual antigens of cattle. Bovine CD8 (BoCD8). Vet Immunol Immunopathol. 27 (1-3): 65-9. 2. Gutierrez, M. et al. (1999) The detection of CD2+, CD4+, CD8+, and WC1+ T lymphocytes, B cells and macrophages in fixed and paraffin embedded bovine tissue using a range of antigen recovery and signal amplification techniques. Vet Immunol Immunopathol. 71 (3-4): 321-34. 3. Winkler, M.T. et al. (1999) Bovine herpesvirus 1 can infect CD4(+) T lymphocytes and induce programmed cell death during acute infection of cattle. J Virol. 73 (10): 8657-68. 4. Winkler, M.T. et al. (2000) Persistence and reactivation of bovine herpesvirus 1 in the tonsils of latently infected calves. J Virol. 74 (11): 5337-46. 5. Twizere, J.C. et al. (2000) Discordance between bovine leukemia virus tax immortalization <i>in vitro</i> and oncogenicity <i>in vivo</i>. J Virol. 74 (21): 9895-902. 6. Harris, J. et al. (2002) Expression of caveolin by bovine lymphocytes and antigen-presenting cells. Immunology. 105: 190-5. 7. Toman, M. et al. (2003) Immunological characteristics of cattle with <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> infection Vet. Med. – Czech, 48, 2003: 147–54. 8. Vordermeier, H.M. et al. (2004) Cellular immune responses induced in cattle by heterologous prime-boost vaccination using recombinant viruses and bacille Calmette-Guérin. Immunology. 112: 461-70. 9. Vitale, F. et al. (2006) ESAT-6 peptide recognition by bovine CD8+ lymphocytes of naturally infected cows in herds from southern Italy. Clin Vaccine Immunol. 13: 530-3.

10. Fulton, B.E. Jr. *et al.* (2006) Dissemination of bovine leukemia virus-infected cells from a newly infected sheep lymph node. [J Virol. 80: 7873-84.](#)
11. Liebana, E. *et al.* (2007) Distribution and activation of T-lymphocyte subsets in tuberculous bovine lymph-node granulomas. [Vet Pathol. 44: 366-72.](#)
12. Foulon, E. & Foucras, G. (2008) Two populations of ovine bone marrow-derived dendritic cells can be generated with recombinant GM-CSF and separated on CD11b expression. [J Immunol Methods. 339 \(1\): 1-10.](#)
13. Sidders, B. *et al.* (2008) Screening of highly expressed mycobacterial genes identifies Rv3615c as a useful differential diagnostic antigen for the *Mycobacterium tuberculosis* complex. [Infect Immun. 76: 3932-9.](#)
14. Lynch, E.M. *et al.* (2010) Effect of abrupt weaning at housing on leukocyte distribution, functional activity of neutrophils, and acute phase protein response of beef calves. [BMC Vet Res. 6: 39.](#)
15. Coad, M. *et al.* (2010) Repeat tuberculin skin testing leads to desensitisation in naturally infected tuberculous cattle which is associated with elevated interleukin-10 and decreased interleukin-1 beta responses. [Vet Res. 41: 14.](#)
16. Constantinoiu, C.C. *et al.* (2010) Local immune response against larvae of *Rhipicephalus (Boophilus) microplus* in *Bos taurus indicus* and *Bos taurus taurus* cattle. [Int J Parasitol. 40: 865-75.](#)
17. La Manna, M.P. *et al.* (2011) Expansion of intracellular IFN- γ positive lymphocytes during *Mycoplasma agalactiae* infection in sheep. [Res Vet Sci. 91 \(3\): e64-7.](#)
18. Sanchez, J. *et al.* (2011) Microscopical and immunological features of tuberculoid granulomata and cavitory pulmonary tuberculosis in naturally infected goats. [J Comp Pathol. 145 \(2-3\): 107-17.](#)
19. Lacroux, C. *et al.* (2012) Prionemia and leukocyte-platelet-associated infectivity in sheep transmissible spongiform encephalopathy models. [J Virol. 86 \(4\): 2056-66.](#)
20. Brodzki, P. *et al.* (2014) Phenotyping of leukocytes and granulocyte and monocyte phagocytic activity in the peripheral blood and uterus of cows with endometritis. [Theriogenology. 82 \(3\): 403-10.](#)
21. Silva, A.P. *et al.* (2015) Encapsulated *Brucella ovis* Lacking a Putative ATP-Binding Cassette Transporter (ΔabcBA) Protects against Wild Type *Brucella ovis* in Rams. [PLoS One. 10 \(8\): e0136865.](#)
22. Leite FL *et al.* (2015) ZAP-70, CTLA-4 and proximal T cell receptor signaling in cows infected with *Mycobacterium avium* subsp. *paratuberculosis*. [Vet Immunol Immunopathol. 167 \(1-2\): 15-21.](#)
23. Stenfeldt, C. *et al.* (2015) Pathogenesis of Primary Foot-and-Mouth Disease Virus Infection in the Nasopharynx of Vaccinated and Non-Vaccinated Cattle. [PLoS One. 10 \(11\): e0143666.](#)
24. Romero-Palomo, F. *et al.* (2017) Immunopathologic Changes in the Thymus of Calves Pre-infected with BVDV and Challenged with BHV-1. [Transbound Emerg Dis. 64 \(2\): 574-84.](#)
25. Schmidt, N. *et al.* (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant *Escherichia coli* Shiga toxins. [Vet Res. 49 \(1\): 28.](#)
26. Pérez-caballero, R. *et al.* (2018) Comparative dynamics of peritoneal cell immunophenotypes in sheep during the early and late stages of the infection with *Fasciola hepatica* by flow cytometric analysis. [Parasit Vectors. 11 \(1\): 640.](#)

27. Benedictus, L. *et al.* (2019) Immunization of young heifers with staphylococcal immune evasion proteins before natural exposure to *Staphylococcus aureus* induces a humoral immune response in serum and milk. [BMC Vet Res. 15 \(1\): 15.](#)
28. Nakajima, N. *et al.* (2019) Effects of direct exposure to cold weather under grazing in winter on the physiological, immunological, and behavioral conditions of Japanese Black beef cattle in central Japan. [Anim Sci J. 90 \(8\): 1033-41.](#)
29. de Araújo, F.F. *et al.* (2019) Distinct immune response profile during *Rhipicephalus (Boophilus) microplus*. infestations of guzerat dairy herd according to the maternal lineage ancestry (mitochondrial DNA). [Vet Parasitol. 273: 36-44.](#)
30. Kolar, Q.K. *et al.* (2020) Anatomical distribution of respiratory tract leukocyte cell subsets in neonatal calves. [Vet Immunol Immunopathol. 227: 110090.](#)
31. Risalde, M.A. *et al.* (2020) BVDV permissiveness and lack of expression of co-stimulatory molecules on PBMCs from calves pre-infected with BVDV. [Comp Immunol Microbiol Infect Dis. 68: 101388.](#)
32. Bidart, J. *et al.* (2020) A New Cage-Like Particle Adjuvant Enhances Protection of Foot-and-Mouth Disease Vaccine. [Front Vet Sci. 7: 396.](#)
33. Brodzki, P. *et al.* (2020) Selected leukocyte subpopulations in peripheral blood and uterine washings in cows before and after intrauterine administration of cefapirin and methisoprinol. [Anim Sci J. 91 \(1\): e13306.](#)
34. Bloomer, S.A. *et al.* (2020) Aging results in accumulation of M1 and M2 hepatic macrophages and a differential response to gadolinium chloride. [Histochem Cell Biol. 153 \(1\): 37-48.](#)
35. Gondaira, S. *et al.* (2020) Immunosuppression in Cows following Intramammary Infusion of *Mycoplasma bovis*. [Infect Immun. 88 \(3\) :e00521-19.](#)
36. Damani-Yokota, P. *et al.* (2021) Transcriptional programming and gene regulation in WC1⁺ γδ T cell subpopulations. [Mol Immunol. 142: 50-62.](#)
37. Sirak, A. *et al.* (2021) Cellular and Cytokine Responses in Lymph Node Granulomas of *Bacillus Calmette Guérin* (BCG)-Vaccinated and Non-vaccinated Cross-Breed Calves Naturally Infected With *Mycobacterium bovis*. [Front Vet Sci. 8: 698800.](#)
38. Colombatti, M.O. *et al.* (2021) Evaluation of a virulent strain of *Mycobacterium avium* subsp. *paratuberculosis* used as a heat-killed vaccine. [Vaccine. 39 \(51\): 7401-12.](#)
39. Park, D.S. *et al.* (2021) Dynamic changes in blood immune cell composition and function in Holstein and Jersey steers in response to heat stress. [Cell Stress Chaperones. 26 \(4\): 705-20.](#)
40. Nashiruddullah, N. *et al.* (2021) Dermal Response to Experimental Orf virus (ORFV) Infection in Goats, Mice and Rabbit [Indian J Anim Res. 56 \(8\): B-4266 1003-9.](#)
41. Kato-Mori, Y. *et al.* (2021) Characterization of a variant CD4 molecule in Japanese Black cattle. [Vet Immunol Immunopathol. 232: 110167.](#)
42. Casaro, S. *et al.* (2022) Flow cytometry panels for immunophenotyping dairy cattle peripheral blood leukocytes [Vet Immunol Immunopathol. 248: 110417.](#)
43. Elsayed, M.S.A.E. *et al.* (2022) Real-time PCR using *atpE*, conventional PCR targeting different regions of difference, and flow cytometry for confirmation of *Mycobacterium bovis*. in buffaloes and cattle from the Delta area of Egypt. [BMC Microbiol. 22 \(1\): 154.](#)
44. Korbonits, L. *et al.* (2022) *Mycobacterium avium* subsp. *paratuberculosis* Infected Cows Reveal Divergent Immune Response in Bovine Peripheral Blood Derived Lymphocyte Proteome. [Metabolites. 12 \(10\): 924.](#)

45. Tucker, N. *et al.* (2023) Bovine blood and milk T-cell subsets in distinct states of activation and differentiation during subclinical *Staphylococcus aureus* mastitis. [J Reprod Immunol. 156: 103826.](#)
46. Özbek, M. & Bayraktaroğlu, A.G. (2019) Developmental study on the ileal Peyer's patches of sheep, and cytokeratin-18 as a possible marker for M cells in follicle associated epithelium. [Acta Histochem. 121 \(3\): 311-22.](#)
47. Benedictus, L. *et al.* (2019) Immunization of young heifers with staphylococcal immune evasion proteins before natural exposure to *Staphylococcus aureus* induces a humoral immune response in serum and milk. [BMC Vet Res. 15 \(1\): 15.](#)
48. Yang, L. *et al.* (2018) Association of the expression of Th cytokines with peripheral CD4 and CD8 lymphocyte subsets after vaccination with FMD vaccine in Holstein young sires. [Res Vet Sci. 119: 79-84.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA837GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#), [DyLight@650](#), [DyLight@680](#), [DyLight@800](#), [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
- Goat Anti Mouse IgG (STAR77...) [HRP](#)
- Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

'M381498:210512'

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 16 Feb 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)