

Datasheet: MCA87 BATCH NUMBER 167886

Description:	escription: MOUSE ANTI HUMAN CD4	
Specificity:	CD45	
Other names:	LCA	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	F10-89-4	
Isotype:	lgG2a	
Quantity:	0.2 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/10 - 1/50
Immunohistology - Frozen	-			1/500 - 1/1000
Immunohistology - Paraffin (1)	-			1/100 - 1/200
ELISA				
Immunoprecipitation	-			
Western Blotting			•	
Rare Cell / CTC				1/200
Enumeration	-			1/200

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.

EDTA pH8.0 is recommended for this purpose.

Target Species	Human
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human T lymphocytes.
External Database Links	UniProt: P08575 Related reagents Entrez Gene: 5788 PTPRC Related reagents
Synonyms	CD45
RRID	AB_871979
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	Mouse anti Human CD45 antibody, clone F10-89-4 recognizes the human CD45 cell surface antigen, also known as leucocyte common antigen (LCA). CD45 is a complex molecule existing in a number of isoforms.
	Antibodies recognizing a common epitope on all of these isoforms are termed CD45 whilst those recognizing only individual isoforms are termed CD45RA or CD45RO etc.
	Mouse anti Human CD45 antibody, clone F10-89-4 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes, having a higher level of expression on lymphocytes than on granulocytes. It is routinely tested in flow cytometry on human peripheral blood leucocytes.
	Mouse anti Human CD45 antibody, clone F10-89-4, has been validated for use on the Genesis Cell Isolation System with the CelSelect Slide TM technology.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human tonsil

References

- 1. Quenby, S *et al.* (1999) Pre-implantation endometrial leukocytes in women with recurrent miscarriage. Human Reprod. 14(9):2386-2391.
- 2. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. <u>Am J Pathol. 177: 2011-21.</u>
- 3. Mallam, E. *et al.* (2010) Characterization of *in vitro* expanded bone marrow-derived mesenchymal stem cells from patients with multiple sclerosis. Mult Scler. 16: 909-18.
- 4. Marrinucci, D. *et al.* (2010) Cytomorphology of circulating colorectal tumor cells:a small case series. <u>J Oncol. 2010: 861341.</u>
- 5. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. PLoS One. 7: e35577.
- 6. De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. <u>Cytometry A. 81 (4): 312-23.</u>
- 7. Kazane, S.A. *et al.* (2012) Site-specific DNA-antibody conjugates for specific and sensitive immuno-PCR. Proc Natl Acad Sci U S A. 109: 3731-6.
- 8. Spaas, J.H. *et al.* (2013) Culture and characterisation of equine peripheral blood mesenchymal stromal cells. Vet J. 195 (1): 107-13.
- 9. Sadarangani, A. *et al.* (2015) GLI2 inhibition abrogates human leukemia stem cell dormancy. <u>J Transl Med. 13: 98.</u>
- 10. Gunawardene, P. *et al.* (2015) Association Between Circulating Osteogenic Progenitor Cells and Disability and Frailty in Older Persons: The Nepean Osteoporosis and Frailty Study. <u>J Gerontol A Biol Sci Med Sci. pii: glv190.</u>
- 11. Mohamed Suhaimi, N.A. *et al.* (2015) Non-invasive sensitive detection of KRAS and BRAF mutation in circulating tumor cells of colorectal cancer patients. <u>Mol Oncol. 9 (4):</u> 850-60.
- 12. Ruiz, C. *et al.* (2015) Limited genomic heterogeneity of circulating melanoma cells in advanced stage patients. <u>Phys Biol. 12 (1): 016008.</u>
- 13. Gogoi P *et al.* (2016) Development of an Automated and Sensitive Microfluidic Device for Capturing and Characterizing Circulating Tumor Cells (CTCs) from Clinical Blood Samples. PLoS One. 11 (1): e0147400.
- 14. Gomiero, C. *et al.* (2016) Tenogenic induction of equine mesenchymal stem cells by means of growth factors and low-level laser technology. Vet Res Commun. 40 (1): 39-48.
- 15. Bianchessi, M. *et al.* (2016) Effect of Fibroblast Growth Factor 2 on Equine Synovial Fluid Chondroprogenitor Expansion and Chondrogenesis. <u>Stem Cells Int. 2016: 9364974.</u>
- 16. Branly, T. *et al.* (2017) Characterization and use of Equine Bone Marrow Mesenchymal Stem Cells in Equine Cartilage Engineering. Study of their Hyaline Cartilage Forming Potential when Cultured under Hypoxia within a Biomaterial in the Presence of BMP-2 and TGF-β1. Stem Cell Rev Rep. 13 (5): 611-30.
- 17. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. <u>PLoS One</u>. 13 (2): e0192244.
- 18. Shishido, S.N. *et al.* (2019) Circulating tumor cells as a response monitor in stage IV non-small cell lung cancer. <u>J Transl Med. 17 (1): 294.</u>
- 19. Welter, L. *et al.* (2020) Treatment response and tumor evolution: lessons from an extended series of multianalyte liquid biopsies in a metastatic breast cancer patient. <u>Cold Spring Harb Mol Case Stud. 6 (6): a005819.</u>
- 20. Ndacayisaba, L.J. *et al.* (2022) Enrichment-Free Single-Cell Detection and Morphogenomic Profiling of Myeloma Patient Samples to Delineate Circulating Rare Plasma Cell Clones <u>Curr Oncol. 29 (5): 2954-72.</u>

- 21. Shishido, S.N. *et al.* (2022) Liquid Biopsy Landscape in Patients with Primary Upper Tract Urothelial Carcinoma. Cancers (Basel). 14 (12): 3007.
- 22. Chai, S. *et al.* (2022) Identification of epithelial and mesenchymal circulating tumor cells in clonal lineage of an aggressive prostate cancer case. NPJ Precis Oncol. 6 (1): 41.
- 23. Zhu, J. *et al.* (2022) Sequential Method for Analysis of CTCs and Exosomes from the Same Sample of Patient Blood. ACS Omega. 7 (42): 37581-88.
- 24. Setayesh, S.M. *et al.* (2022) Multianalyte liquid biopsy to aid the diagnostic workup of breast cancer. NPJ Breast Cancer. 8 (1): 112.
- 25. Ndacayisaba, L.J. *et al.* (2022) Characterization of BCMA Expression in Circulating Rare Single Cells of Patients with Plasma Cell Neoplasms. <u>Int J Mol Sci. 23 (21): 13427.</u>
- 26. Qi, E. *et al.* (2023) Investigation of liquid biopsy analytes in peripheral blood of individuals after SARS-CoV-2 infection. <u>EBioMedicine</u>. 90: 104519.
- 27. Seo, J. *et al.* (2023) Plasticity of circulating tumor cells in small cell lung cancer. <u>Sci</u> Rep. 13 (1): 11775.
- 28. Setayesh, S.M. *et al.* (2023) Targeted single-cell proteomic analysis identifies new liquid biopsy biomarkers associated with multiple myeloma. NPJ Precis Oncol. 7 (1): 95.
- 29. Welter, L. *et al.* (2023) Cell State and Cell Type: Deconvoluting Circulating Tumor Cell Populations in Liquid Biopsies by Multi-Omics. Cancers (Basel). 15 (15): 3949.
- 30. Shishido, S.N. *et al.* (2024) Cancer-related cells and oncosomes in the liquid biopsy of pancreatic cancer patients undergoing surgery. <u>NPJ Precis Oncol. 8 (1): 36.</u>

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Acknowledgements	CelSelect Slides is a trademark of Bio-Rad Laboratories, Inc. in certain jurisdictions.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA87 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426154:231121'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint