Datasheet: MCA87PB BATCH NUMBER 0715

Description:	MOUSE ANTI HUMAN CD45:Pacific Blue®
Specificity:	CD45
Other names:	LCA
Format:	Pacific Blue®
Product Type:	Monoclonal Antibody
Clone:	F10-89-4
Isotype:	lgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications	derived from testing wi communications from t information. For gener	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .			
		Yes No	Not Determined	Suggested Dilution	
	Flow Cytometry	•		Neat	
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.				
Target Species	Human				
Species Cross	Reacts with: Horse				
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG conjugated to Pacific Blue® - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	Pacific Blue®	410	455		
Preparation	Purified IgG prepared supernatant	by affinity chromatog	raphy on Protein A fror	n tissue culture	

Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin	
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml	
Immunogen	Human T lymphocytes.	
External Database Links	UniProt: P08575 Related reagents Entrez Gene: 5788 PTPRC Related reagents	
Synonyms	CD45	
RRID	AB_566767	
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells o myeloma cell line.	f the mouse NS-1
Specificity	Mouse anti Human CD45 antibody, clone F10-89-4 recognizes to surface antigen, also known as the leucocyte common antigen (LC molecule existing in a number of isoforms.	CA). CD45 is a complex
	Antibodies recognising a common epitope on all of these isoforms those recognising only individual isoforms are termed CD45RA or	
	Mouse anti Human CD45 antibody, clone F10-89-4 reacts with all expressed by all haematopoietic cells, except erythrocytes, having expression on lymphocytes than on granulocytes.	
	Mouse anti Human CD45 antibody, clone F10-89-4 is routinely tes human peripheral blood leucocytes	ted in flow cytometry on
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100	Oul whole blood.
References	 Dalchau, R. <i>et al.</i> (1980) Monoclonal antibody to a human leuko glycoprotein probably homologous to the leukocyte-common (L-C) <u>Immunol. 10 (10): 737-44.</u> Quenby, S <i>et al.</i> (1999) Pre-implantation endometrial leukocytes recurrent miscarriage. <u>Human Reprod. 14(9):2386-2391.</u> Hauser, P.V. <i>et al.</i> (2010) Stem cells derived from human amnio acute kidney injury recovery. <u>Am J Pathol. 177: 2011-21.</u> Mallam, E. <i>et al.</i> (2010) Characterization of <i>in vitro</i> expanded box 	antigen of the rat. <u>Eur J</u> s in women with tic fluid contribute to

mesenchymal stem cells from patients with multiple sclerosis. Mult Scler. 16: 909-18.

5. Marrinucci, D. *et al.* (2010) Cytomorphology of circulating colorectal tumor cells:a small case series. <u>J Oncol. 2010: 861341.</u>

6. Kazane, S.A. *et al.* (2012) Site-specific DNA-antibody conjugates for specific and sensitive immuno-PCR. <u>Proc Natl Acad Sci U S A. 109: 3731-6.</u>

7. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. <u>PLoS One. 7: e35577.</u>

8. Sadarangani, A. *et al.* (2015) GLI2 inhibition abrogates human leukemia stem cell dormancy. <u>J Transl Med. 13: 98.</u>

9. Gunawardene, P. *et al.* (2015) Association Between Circulating Osteogenic Progenitor Cells and Disability and Frailty in Older Persons: The Nepean Osteoporosis and Frailty Study. J Gerontol A Biol Sci Med Sci. pii: glv190.

10. Gogoi P *et al.* (2016) Development of an Automated and Sensitive Microfluidic Device for Capturing and Characterizing Circulating Tumor Cells (CTCs) from Clinical Blood Samples. <u>PLoS One. 11 (1): e0147400.</u>

11. Spaas, J.H. *et al.* (2013) Culture and characterisation of equine peripheral blood mesenchymal stromal cells. <u>Vet J. 195 (1): 107-13.</u>

 Gomiero, C. *et al.* (2016) Tenogenic induction of equine mesenchymal stem cells by means of growth factors and low-level laser technology. <u>Vet Res Commun. 40 (1): 39-48.</u>
 De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. <u>Cytometry A. 81 (4):</u> <u>312-23.</u>

Bianchessi, M. *et al.* (2016) Effect of Fibroblast Growth Factor 2 on Equine Synovial Fluid Chondroprogenitor Expansion and Chondrogenesis. <u>Stem Cells Int. 2016: 9364974.</u>
 Mohamed Suhaimi, N.A. *et al.* (2015) Non-invasive sensitive detection of KRAS and BRAF mutation in circulating tumor cells of colorectal cancer patients. <u>Mol Oncol. 9 (4):</u> 850-60.

16. Ruiz, C. *et al.* (2015) Limited genomic heterogeneity of circulating melanoma cells in advanced stage patients. <u>Phys Biol. 12 (1): 016008.</u>

17. Branly, T. *et al.* (2017) Characterization and use of Equine Bone Marrow Mesenchymal Stem Cells in Equine Cartilage Engineering. Study of their Hyaline Cartilage Forming Potential when Cultured under Hypoxia within a Biomaterial in the Presence of BMP-2 and TGF-β1. <u>Stem Cell Rev. Jun 09 [Epub ahead of print].</u>

18. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. <u>PLoS One. 13 (2): e0192244.</u>

19. Shishido, S.N. *et al.* (2019) Circulating tumor cells as a response monitor in stage IV non-small cell lung cancer. <u>J Transl Med. 17 (1): 294.</u>

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	18 months from date of despatch.
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA87PB
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:Pacific Blue® (MCA929PB)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M342017:190110'

Printed on 09 Jul 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint