

## Datasheet: MCA87T

**BATCH NUMBER 1801**

<b>Description:</b>	MOUSE ANTI HUMAN CD45
<b>Specificity:</b>	CD45
<b>Other names:</b>	LCA
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	F10-89-4
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			1/500 - 1/1000
Immunohistology - Paraffin (1)	▪			1/100 - 1/200
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.**

**EDTA pH8.0 is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Horse <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for

further information.

---

<b>Product Form</b>	Purified IgG - liquid
---------------------	-----------------------

---

<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G
--------------------	---

---

<b>Buffer Solution</b>	Phosphate buffered saline
------------------------	---------------------------

---

<b>Preservative Stabilisers</b>	0.09% Sodium Azide
---------------------------------	--------------------

---

<b>Carrier Free</b>	Yes
---------------------	-----

---

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
---------------------------------------	-----------------------------

---

<b>Immunogen</b>	Human T lymphocytes.
------------------	----------------------

---

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P08575</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">5788</a> PTPRC <a href="#">Related reagents</a>
--------------------------------	--

---

<b>Synonyms</b>	CD45
-----------------	------

---

<b>RRID</b>	AB_1102051
-------------	------------

---

<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
------------------------	--

---

<b>Specificity</b>	<p><b>Mouse anti Human CD45 antibody, clone F10-89-4</b> recognizes the human CD45 cell surface antigen, also known as the leucocyte common antigen (LCA). CD45 is a complex molecule existing in a number of isoforms.</p> <p>Antibodies recognising a common epitope on all of these isoforms are termed CD45 whilst those recognising only individual isoforms are termed CD45RA or CD45RO etc.</p> <p>Mouse anti Human CD45 antibody, clone F10-89-4 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes, having a higher level of expression on lymphocytes than on granulocytes.</p> <p>Mouse anti Human CD45 antibody, clone F10-89-4 is routinely tested in flow cytometry on human peripheral blood leucocytes</p>
--------------------	--

---

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
-----------------------	---

---

<b>Histology Positive</b>	Human tonsil
---------------------------	--------------

## References

1. Dalchau, R. *et al.* (1980) Monoclonal antibody to a human leukocyte-specific membrane glycoprotein probably homologous to the leukocyte-common (L-C) antigen of the rat. [Eur J Immunol. 10 \(10\): 737-44.](#)
2. Quenby, S *et al.* (1999) Pre-implantation endometrial leukocytes in women with recurrent miscarriage. [Human Reprod. 14\(9\):2386-2391.](#)
3. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. [Am J Pathol. 177: 2011-21.](#)
4. Mallam, E. *et al.* (2010) Characterization of *in vitro* expanded bone marrow-derived mesenchymal stem cells from patients with multiple sclerosis. [Mult Scler. 16: 909-18.](#)
5. Marrinucci, D. *et al.* (2010) Cytomorphology of circulating colorectal tumor cells:a small case series. [J Oncol. 2010: 861341.](#)
6. Kazane, S.A. *et al.* (2012) Site-specific DNA-antibody conjugates for specific and sensitive immuno-PCR. [Proc Natl Acad Sci U S A. 109: 3731-6.](#)
7. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. [PLoS One. 7: e35577.](#)
8. Sadarangani, A. *et al.* (2015) GLI2 inhibition abrogates human leukemia stem cell dormancy. [J Transl Med. 13: 98.](#)
9. Gunawardene, P. *et al.* (2015) Association Between Circulating Osteogenic Progenitor Cells and Disability and Frailty in Older Persons: The Nepean Osteoporosis and Frailty Study. [J Gerontol A Biol Sci Med Sci. pii: glv190.](#)
10. Gogoi P *et al.* (2016) Development of an Automated and Sensitive Microfluidic Device for Capturing and Characterizing Circulating Tumor Cells (CTCs) from Clinical Blood Samples. [PLoS One. 11 \(1\): e0147400.](#)
11. Spaas, J.H. *et al.* (2013) Culture and characterisation of equine peripheral blood mesenchymal stromal cells. [Vet J. 195 \(1\): 107-13.](#)
12. Gomiero, C. *et al.* (2016) Tenogenic induction of equine mesenchymal stem cells by means of growth factors and low-level laser technology. [Vet Res Commun. 40 \(1\): 39-48.](#)
13. De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. [Cytometry A. 81 \(4\): 312-23.](#)
14. Bianchessi, M. *et al.* (2016) Effect of Fibroblast Growth Factor 2 on Equine Synovial Fluid Chondroprogenitor Expansion and Chondrogenesis. [Stem Cells Int. 2016: 9364974.](#)
15. Mohamed Suhaimi, N.A. *et al.* (2015) Non-invasive sensitive detection of KRAS and BRAF mutation in circulating tumor cells of colorectal cancer patients. [Mol Oncol. 9 \(4\): 850-60.](#)
16. Ruiz, C. *et al.* (2015) Limited genomic heterogeneity of circulating melanoma cells in advanced stage patients. [Phys Biol. 12 \(1\): 016008.](#)
17. Branly, T. *et al.* (2017) Characterization and use of Equine Bone Marrow Mesenchymal Stem Cells in Equine Cartilage Engineering. Study of their Hyaline Cartilage Forming Potential when Cultured under Hypoxia within a Biomaterial in the Presence of BMP-2 and TGF- $\beta$ 1. [Stem Cell Rev. Jun 09 \[Epub ahead of print\].](#)
18. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)
19. Shishido, S.N. *et al.* (2019) Circulating tumor cells as a response monitor in stage IV non-small cell lung cancer. [J Transl Med. 17 \(1\): 294.](#)

20. Welter, L. *et al.* (2020) Treatment response and tumor evolution: Lessons from an extended series of multi-analyte liquid biopsies in a metastatic breast cancer patient. [Cold Spring Harb Mol Case Stud. Nov 17 \[Epub ahead of print\].](#)

---

**Storage** Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA87T>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M369081:200529'

Printed on 07 May 2024