

## Datasheet: MCA88

<b>Description:</b>	MOUSE ANTI HUMAN CD45RA
<b>Specificity:</b>	CD45RA
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	F8-11-13
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			1/200
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Rhesus Monkey</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human T lymphocytes
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P08575</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">5788</a>    PTPRC    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CD45
<b>RRID</b>	AB_871980
<b>Fusion Partners</b>	Spleen cells from immunized BALB/c mouse were fused with cells of the NS-1 mouse myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human CD45RA antibody, clone F8-11-13</b> recognizes the human CD45RA cell surface antigen.</p> <p>CD45 is a complex molecule existing in a number of isoforms. Antibodies recognizing a common epitope on all of these isoforms are termed CD45 whilst those recognizing only individual isoforms are termed CD45RA or CD45RO etc.</p> <p>Mouse anti Human CD45RA antibody, clone F8-11-13 reacts with the high molecular weight form of the leucocyte common antigen (LCA) expressed by B lymphocytes and a subset of T lymphocytes.</p> <p>Mouse anti Human CD45RA antibody, clone F8-11-13 is routinely tested in flow cytometry on human peripheral blood lymphocytes.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $10^6$ cells in 100ul.
<b>Immunohistology</b>	This product does not require protein digestion pre-treatment of paraffin sections. This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections.
<b>Histology Positive Control Tissue</b>	Human tonsil
<b>References</b>	1. Dalchau, R. & Fabre, J.W. (1981) Identification with a monoclonal antibody of a predominantly B lymphocyte-specific determinant of the human leukocyte common

- antigen. Evidence for structural and possible functional diversity of the human leukocyte common molecule. [J Exp Med. 153 \(4\): 753-65.](#)
2. O'Grady, J.T. *et al.* (1994) CD40 expression in Hodgkin's disease. [Am J Pathol. 144 \(1\): 21-6.](#)
3. Franitza, S. *et al.* (2002) TGF-beta1 enhances SDF-1alpha-induced chemotaxis and homing of naive T cells by up-regulating CXCR4 expression and downstream cytoskeletal effector molecules. [Eur J Immunol. 32 \(1\): 193-202.](#)
4. Leigh, J.E. *et al.* (2006) Characterization of the immune status of CD8+ T cells in oral lesions of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. [Clin Vaccine Immunol. 13 \(6\): 678-83.](#)
5. Hunt, L. *et al.* (2016) T cell subsets: an immunological biomarker to predict progression to clinical arthritis in ACPA-positive individuals. [Ann Rheum Dis. 75 \(10\): 1884-9.](#)
6. Ibane, J.A. *et al.* (2012) The major CD8 T cell effector memory subset in the normal and *Chlamydia trachomatis*-infected human endocervix is low in perforin. [BMC Immunol. 13: 66.](#)
7. Ponchel, F. *et al.* (2014) An immunological biomarker to predict MTX response in early RA. [Ann Rheum Dis. 73 \(11\): 2047-53.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA88>  
10040

---

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

- |   |   |
|---|---|
| Goat Anti Mouse IgG (STAR77...)         | <a href="#">HRP</a>   |
| Rabbit Anti Mouse IgG (STAR12...)       | <a href="#">RPE</a>   |
| Goat Anti Mouse IgG (STAR70...)         | <a href="#">FITC</a>  |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | <a href="#">Alk. Phos.</a> , <a href="#">HRP</a>  |
| Rabbit Anti Mouse IgG (STAR9...)        | <a href="#">FITC</a>  |
| Goat Anti Mouse IgG (STAR76...)         | <a href="#">RPE</a>   |
| Goat Anti Mouse IgG (Fc) (STAR120...)   | <a href="#">FITC</a> , <a href="#">HRP</a>  |
| Goat Anti Mouse IgG (H/L) (STAR117...)  | <a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@550</a> ,<br><a href="#">DyLight@650</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> ,<br><a href="#">FITC</a> , <a href="#">HRP</a> |
| Rabbit Anti Mouse IgG (STAR13...)       | <a href="#">HRP</a>   |

## Recommended Negative Controls

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M381519:210512'

**Printed on 25 Mar 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)